


MBP Antibody [clone MBP/4272] (V8676)


| Catalog No. | Formulation | Size |
|----------------|--|--------|
| V8676-100UG | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide | 100 ug |
| V8676-20UG | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide | 20 ug |
| V8676SAF-100UG | 1 mg/ml in 1X PBS; BSA free, sodium azide free | 100 ug |

Bulk quote request


| | |
|---------------------------|--|
| Availability | 1-3 business days |
| Species Reactivity | Human |
| Format | Purified |
| Host | Mouse |
| Clonality | Monoclonal (mouse origin) |
| Isotype | Mouse IgG1 |
| Clone Name | MBP/4272 |
| Purity | Protein G affinity chromatography |
| UniProt | P02686 |
| Localization | Cell surface, cytoplasm |
| Applications | Western Blot : 1-2ug/ml Immunohistochemistry (FFPE) : 1-2ug/ml for 30 minutes at RT |
| Limitations | This MBP antibody is available for research use only. |




IHC staining of FFPE human brain with MBP antibody. HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.




IHC staining of FFPE human brain with MBP antibody. HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



Western blot testing of human brain lysate with MBP antibody. Isoforms may be visualized from 20~37 kDa.



Western blot testing of human U-87 MG cell lysate with MBP antibody. Isoforms may be visualized from 20~37 kDa.



SDS-PAGE analysis of purified, BSA-free MBP antibody as confirmation of integrity and purity.

Description

Myelin basic protein (MBP) is the second most abundant protein in central nervous system (CNS) myelin: it comprises 30% of the total protein and about 10% of the dry weight of myelin. It is the only structural protein found so far to be essential for formation of CNS myelin, and has been called the executive molecule of myelin. MBP can interact with a number of polyanionic proteins including actin, tubulin, calmodulin, and clathrin, and negatively charged lipids, and acquires structure on binding to them. It may act as a membrane actin-binding protein, which might allow it to participate in transmission of extracellular signals to the cytoskeleton in oligodendrocytes and tight junctions in myelin. MBP may be applicable as a marker for oligodendrogiomas. MBP/4272 recognizes an epitope in the 119-131 region of MBP, useful in clinical diagnosis to detect MBP levels in human, rat and cow MBP.

Application Notes

Optimal dilution of the MBP antibody should be determined by the researcher.

Immunogen

Amino acids GAEGQRPGFGYGG were used as the immunogen for the MBP antibody.

Storage

Store the MBP antibody at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).