
MCAM Antibody MCAM/1101 / CD146 Mouse Monoclonal [clone MCAM/1101] (V2696)


Catalog No.	Formulation	Size
V2696-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V2696-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V2696SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug
V2696IHC-7ML	Prediluted in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide; *For IHC use only*	7 ml

Bulk quote request

Species Reactivity	Human
Format	Purified
Host	Mouse
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG1, kappa
Clone Name	MCAM/1101
Purity	Protein G affinity chromatography
UniProt	P43121
Localization	Cell surface, cytoplasm
Applications	Flow cytometry: 0.5-1ug/10^6 cells Immunofluorescence: 0.5-1ug/ml Immunohistochemistry (FFPE): 0.5-1ug/ml for 30 min at RT
Limitations	This MCAM Antibody MCAM/1101 is available for research use only.



MCAM Antibody MCAM/1101 IHC: Formalin-fixed, paraffin-embedded human tonsil tissue stained with MCAM antibody (clone MCAM/1101).



MCAM Antibody MCAM/1101 IHC: Formalin-fixed, paraffin-embedded human melanoma stained with MCAM antibody (MCAM/1101)

Description

The human Mel-CAM gene maps to chromosome 11q23 and encodes a trans-membrane glycoprotein, also designated MCAM, MUC 18 or CD146, that belongs to the immunoglobulin superfamily and functions as a Ca2+-independent cell adhesion molecule. MCAM expression is restricted to advanced primary and metastatic melanomas and to cell lines of the neuroectodermal lineage, but not normal melanocytes. It is found on 80% of advanced primary human melanomas and correlates well with development of metastatic disease.

For an alternative antibody targeting the same cell adhesion molecule, see our MCAM Antibody MCAM/3046, a mouse monoclonal antibody featuring human melanoma IHC and protein microarray specificity validation.

Application Notes

Optimal dilution of the MCAM Antibody MCAM/1101 should be determined by the researcher.

1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min
2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.

Immunogen

Recombinant human protein was used as the immunogen for the MCAM Antibody MCAM/1101.

Storage

Store the MCAM Antibody MCAM/1101 at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).

Alternate Names

MCAM antibody, MUC18 antibody, Mel-CAM antibody, Melanoma Cell Adhesion Molecule antibody, Cell Surface Glycoprotein MUC18 antibody, CD146 Antigen antibody, Recombinant MCAM antibody, Recombinant MUC18 antibody

Copyright © NSJ Bioreagents. All rights reserved.