
MCAM Antibody MCAM/3048 / MUC18 Mouse MAb [clone MCAM/3048] (V7942)


Catalog No.	Formulation	Size
V7942-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V7942-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V7942SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

Bulk quote request


Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Host	Mouse
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG2b, kappa
Clone Name	MCAM/3048
Purity	Protein G affinity chromatography
UniProt	P43121
Localization	Cell surface, cytoplasmic
Applications	Immunohistochemistry (FFPE): 1-2ug/ml
Limitations	This MCAM Antibody MCAM/3048 is available for research use only.



IHC staining of FFPE human melanoma with MCAM Antibody MCAM/3048. HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



IHC staining of FFPE human melanoma with MCAM antibody (clone MCAM/3048). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



Analysis of HuProt(TM) microarray containing more than 19,000 full-length human proteins using MCAM antibody (clone MCAM/3048). These results demonstrate the foremost specificity of the MCAM/3048 mAb.
Z- and S- score: The Z-score represents the strength of a signal that an antibody (in combination with a fluorescently-tagged anti-IgG secondary Ab) produces when binding to a particular protein on the HuProt(TM) array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If the targets on the HuProt(TM) are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-scores. The S-score therefore represents the relative target specificity of an Ab to its intended target.

Description

The human Mel-CAM gene maps to chromosome 11q23 and encodes a trans-membrane glycoprotein, also designated MCAM, MUC 18 or CD146, that belongs to the immunoglobulin superfamily and functions as a Ca2+-independent cell adhesion molecule. Mel-CAM expression is restricted to advanced primary and metastatic melanomas and to cell lines of the neuroectodermal lineage, but not normal melanocytes. Mel-CAM is found on 80% of advanced primary human melanomas and correlates well with development of metastatic disease.

For an alternative antibody targeting the same cell adhesion molecule, see our MCAM Antibody MCAM/3046, a mouse monoclonal antibody featuring human melanoma IHC and protein microarray specificity validation.

Application Notes

Optimal dilution of the MCAM Antibody MCAM/3048 should be determined by the researcher.

Immunogen

A recombinant human partial protein (amino acids 226-374) was used as the immunogen for the MCAM Antibody MCAM/3048.

Storage

Store the MCAM Antibody MCAM/3048 at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).

Alternate Names

MCAM antibody, MUC18 antibody, Mel-CAM antibody, Melanoma Cell Adhesion Molecule antibody, Cell Surface Glycoprotein MUC18 antibody, CD146 Antigen antibody, Recombinant MCAM antibody, Recombinant MUC18 antibody

Copyright © NSJ Bioreagents. All rights reserved.