


MEF2D Antibody [clone PCRP-MEF2D-3A4] (V9704)


Catalog No.	Formulation	Size
V9704-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	100 ug
V9704-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	20 ug
V9704SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

[Bulk quote request](#)


Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Host	Mouse
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG2b
Clone Name	PCRP-MEF2D-3A4
Purity	Protein A/G affinity
UniProt	Q14814
Localization	Nucleus
Applications	Flow Cytometry : 1-2ug/million cells Immunofluorescence : 1-2ug/ml Immunohistochemistry (FFPE) : 1-2ug/ml
Limitations	This MEF2D antibody is available for research use only.




Immunofluorescent staining of PFA-fixed human HeLa cells using MEF2D antibody (green, clone PCRP-MEF2D-3A4) and phalloidin (red).




FACS staining of PFA-fixed human HeLa cells with MEF2D antibody (blue, clone PCRP-MEF2D-3A4) and isotype control (red).



IHC staining of FFPE human rhabdomyosarcoma tissue with MEF2D antibody (clone PCRP-MEF2D-3A4). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



SDS-PAGE analysis of purified, BSA-free MEF2D antibody (clone PCRP-MEF2D-3A4) as confirmation of integrity and purity.



Analysis of HuProt(TM) microarray containing more than 19,000 full-length human proteins using MEF2D antibody (clone PCRP-MEF2D-3A4). These results demonstrate the foremost specificity of the PCRP-MEF2D-3A4 mAb. Z- and S- score: The Z-score represents the strength of a signal that an antibody (in combination with a fluorescently-tagged anti-IgG secondary Ab) produces when binding to a particular protein on the HuProt(TM) array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If the targets on the HuProt(TM) are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-scores. The S-score therefore represents the relative target specificity of an Ab to its intended target.

Description

The myocyte enhancer factor-2 (MEF-2) family of transcription factors associate with co-repressors or co-activators to regulate development and function of T cells, neuronal cells and muscle cells. Four family members arise from alternatively spliced transcripts, termed MEF-2A, -2B, -2C and -2D. These members bind as homo- and heterodimers to the MEF-2 site in the promoter region of affected genes. Differential regulation in the expression of the four transcripts implies functional distinction for each during embryogenesis and development. The process of differentiation from mesodermal precursor cells to myoblasts has led to the discovery of a variety of tissue-specific factors that regulate muscle gene expression. The myogenic basic helix-loop-helix proteins, including MyoD, myogenin, Myf-5 and MRF4, are one class of identified factors. A second family of DNA-binding regulatory proteins is the myocyte-specific enhancer factor-2 (MEF-2) family. Each of these proteins binds to the MEF-2 target DNA sequence present in the regulatory regions of many muscle-specific genes.

Application Notes

Optimal dilution of the MEF2D antibody should be determined by the researcher.

Immunogen

Recombinant full-length human MEF2D protein was used as the immunogen for the MEF2D antibody.

Storage

Aliquot the MEF2D antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.