


# Melan-A Antibody / MART-1 [clone MLANA/8365R] (V5290)

Catalog No.	Formulation	Size
V5290-100UG	0.2~mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	100 ug
V5290-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	20 ug
V5290SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug


## Recombinant RABBIT MONOCLONAL

# **Bulk quote request**

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Clonality	Recombinant Rabbit Monoclonal
Isotype	Rabbit IgG, kappa
Clone Name	MLANA/8365R
Purity	Protein A/G affinity
UniProt	Q16655
Localization	Cytoplasm
Applications	Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT
Limitations	This Melan-A antibody is available for research use only.



IHC staining of FFPE human melanoma tissue with Melan-A antibody (clone MLANA/8365R). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



### **Description**

This antibody recognizes a protein doublet of 20-22kDa, identified as MART-1 (Melanoma Antigen Recognized by T cells 1) or Melan-A. MART-1 is a newly identified melanocyte differentiation antigen recognized by autologous cytotoxic T lymphocytes. Seven other melanoma associated antigens recognized by autologous cytotoxic T cells include MAGE-1, MAGE-3, tyrosinase, gp100, gp75, BAGE-1, and GAGE-1. Subcellular fractionation shows that MART-1 is present in melanosomes and endoplasmic reticulum. This mAb labels melanomas and other tumors showing melanocytic differentiation. It is also a useful positive-marker for angiomyolipomas. It does not stain tumor cells of epithelial, lymphoid, glial, or mesenchymal origin.

#### **Application Notes**

Optimal dilution of the Melan-A antibody should be determined by the researcher.

#### **Immunogen**

A recombinant fragment corresponding to the N-terminus of human Melan A protein was used as the immunogen for the Melan-A antibody.

## **Storage**

Aliquot the Melan-A antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.