


Melanoma gp100 Antibody / PMEL17 [clone MSSG95-2] (V3916)

Catalog No.	Formulation	Size
V3916-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V3916-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V3916SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

[Bulk quote request](#)

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG1, kappa
Clone Name	MSSG95-2
Purity	Protein G affinity chromatography
UniProt	P40967
Localization	Cytoplasmic
Applications	Immunohistochemistry (FFPE) : 1-2ug/ml for 30 min at RT
Limitations	This Melanoma gp100 antibody is available for research use only.



IHC testing of human melanoma with Melanoma gp100 antibody (clone MSSG95-2). Required HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 min.

Description

PMEL17/Melanoma gp100/SILV plays a central role in the biogenesis of melanosomes. Involved in the maturation of melanosomes from stage I to II. The transition from stage I melanosomes to stage II melanosomes involves an elongation

of the vesicle, and the appearance within of distinct fibrillar structures. Release of the soluble form, ME20-S, could protect tumor cells from antibody mediated immunity. [UniProt]

Application Notes

The stated application concentrations are suggested starting points. Titration of the Melanoma gp100 antibody may be required due to differences in protocols and secondary/substrate sensitivity.

Immunogen

A portion of amino acids 376-502 from the human protein was used as the immunogen for the Melanoma gp100 antibody.

Storage

Store the Melanoma gp100 antibody at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).