
MerTK Monoclonal Antibody MERTK/3022 / c-Mer Mouse Monoclonal Antibody [clone MERTK/3022] (V7663)


Catalog No.	Formulation	Size
V7663-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V7663-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V7663SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

Bulk quote request


Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Host	Mouse
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG2a
Clone Name	MERTK/3022
Purity	Protein G affinity chromatography
UniProt	Q12866
Localization	Cell surface
Applications	1-2ug/ml
Limitations	This MerTK Monoclonal Antibody MERTK/3022 / c-Mer Mouse Monoclonal Antibody is available for research use only.




MerTK Monoclonal Antibody MERTK/3022 / c-Mer Mouse Monoclonal Antibody Immunohistochemistry. IHC staining of FFPE human colon carcinoma with MerTK antibody (clone MERTK/3022). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min and allow to cool before testing.



IHC staining of FFPE human colon carcinoma with MerTK antibody (clone MERTK/3022). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min and allow to cool before testing.



Analysis of HuProt(TM) microarray containing more than 19,000 full-length human proteins using MerTK antibody (clone MERTK/3022). These results demonstrate the foremost specificity of the MERTK/3022 mAb.
Z- and S- score: The Z-score represents the strength of a signal that an antibody (in combination with a fluorescently-tagged anti-IgG secondary Ab) produces when binding to a particular protein on the HuProt(TM) array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If the targets on the HuProt(TM) are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-scores. The S-score therefore represents the relative target specificity of an Ab to its intended target.



SDS-PAGE analysis of purified, BSA-free MerTK antibody (clone MERTK/3022) as confirmation of integrity and purity.

Description

MerTK Monoclonal Antibody MERTK/3022 detects MerTK, also called c-Mer, a member of the Mer/Axl/Tyro3 receptor kinase family. It is a 984 residue transmembrane protein made up of one tyrosine kinase domain, two Fibronectin type-III domains and two immunoglobulinlike C2-type domains. MerTK is the mammalian ortholog of the chicken retroviral oncogene product v-Eyk. This protein plays a critical role in macrophage activation, platelet aggregation, clot stability and the efficient removal of apoptotic cells. Specifically, MerTK acts as a signaling molecule, triggering outer segment ingestion in the retinal pigment epithelium (RPE) phagocytic process. Evidence suggests that MerTK signals via interaction with phosphatidylinositol-specific phospholipase C 2). When the gene encoding for MerTK is mutated, the RPE phagocytosis pathway is disrupted and autosomal recessive retinitis pigmentosa (RP) may result, leading to degeneration of retinal photoreceptor cells.

For a Protein Microarray Validated monoclonal antibody to this target, see our MerTK Antibody MERTK/3015 / MERTK Monoclonal Antibody.

Application Notes

Optimal dilution of the MerTK Monoclonal Antibody MERTK/3022 / c-Mer Mouse Monoclonal Antibody should be determined by the researcher.

Immunogen

A recombinant human partial protein (amino acids 55-148) was used as the immunogen for the MerTK Monoclonal Antibody MERTK/3022.

Storage

Store the MerTK Monoclonal Antibody MERTK/3022 at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).

Copyright © NSJ Bioreagents. All rights reserved.