


MERTK Antibody [clone MERTK/3023] (V8905)


| Catalog No. | Formulation | Size |
|----------------|---|--------|
| V8905-100UG | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide | 100 ug |
| V8905-20UG | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide | 20 ug |
| V8905SAF-100UG | 1 mg/ml in 1X PBS; BSA free, sodium azide free | 100 ug |

Bulk quote request

| | |
|--------------------|---|
| Availability | 1-3 business days |
| Species Reactivity | Human |
| Format | Purified |
| Host | Mouse |
| Clonality | Monoclonal (mouse origin) |
| Isotype | Mouse IgG2a |
| Clone Name | MERTK/3023 |
| Purity | Protein A/G affinity |
| UniProt | Q12866 |
| Localization | Cell surface |
| Applications | Immunohistochemistry (FFPE) : 1-2ug/ml |
| Limitations | This MERTK antibody is available for research use only. |




IHC staining of FFPE human colon carcinoma tissue with MERTK antibody (clone MERTK/3023). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



IHC staining of FFPE human colon carcinoma tissue with MERTK antibody (clone MERTK/3023). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.

Human Protein Microarray Specificity Validation



Analysis of HuProt(TM) microarray containing more than 19,000 full-length human proteins using MERTK antibody (clone MERTK/3023). These results demonstrate the foremost specificity of the MERTK/3023 mAb. Z- and S- score: The Z-score represents the strength of a signal that an antibody (in combination with a fluorescently-tagged anti-IgG secondary Ab) produces when binding to a particular protein on the HuProt(TM) array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If the targets on the HuProt(TM) are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-scores. The S-score therefore represents the relative target specificity of an Ab to its intended target.

Description

MERTK, also called c-Mer, is a member of the Mer/Axl/Tyro3 receptor kinase family. It protein plays a critical role in macrophage activation, platelet aggregation, clot stability and the efficient removal of apoptotic cells.

Application Notes

Optimal dilution of the MERTK antibody should be determined by the researcher.

Immunogen

A portion of amino acids 55-148 was used as the immunogen for the MERTK antibody.

Storage

Aliquot the MERTK antibody and store frozen at -20°C or colder. Avoid repeated freeze-thaw cycles.