


MICA Antibody (F51246)


Catalog No.	Formulation	Size
F51246-0.4ML	In 1X PBS, pH 7.4, with 0.09% sodium azide	0.4 ml
F51246-0.08ML	In 1X PBS, pH 7.4, with 0.09% sodium azide	0.08 ml

[Bulk quote request](#)


Availability	1-3 business days
Species Reactivity	Human
Format	Antigen affinity purified
Host	Rabbit
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit Ig
Purity	Antigen affinity
UniProt	Q29983
Applications	Western Blot : 1:1000 Flow Cytometry : 1:25 Immunofluorescence : 1:25 Immunohistochemistry (FFPE) : 1:25
Limitations	This MICA antibody is available for research use only.




Western blot testing of MICA antibody at 1:2000 dilution. Lane 1: Jurkat lysate; 2: Daudi ; 3: A431; 4: MCF-7; 5: U-87MG; 6: THP-1; Predicted molecular weight: ~43 kDa but may be observed at 38-62 kDa depending on truncation and glycosylation level.




Western blot testing of MICA antibody at 1:2000 dilution. Lane 1: U-87 MG lysate; 2: A431; 3: Jurkat; Predicted molecular weight: ~43kDa but may be observed at 38-62kDa depending on truncation and glycosylation level.




Western blot testing of MICA antibody at 1:1000 dilution + HeLa lysate




Flow cytometric analysis of SK-BR-3 cells using MICA antibody (green) compared to an isotype control of [rabbit IgG \(blue\)](#). Ab was diluted at 1:25 dilution. An Alexa Fluor 488 goat anti-rabbit IgG was used as the secondary.



Immunofluorescent staining of fixed and permeabilized human HeLa cells with MICA antibody (green), phalloidin (red, cytoplasmic actin) and DAPI (blue, nuclear).



Immunofluorescent staining of fixed and permeabilized human HeLa cells with MICA antibody (green), phalloidin (red, cytoplasmic actin) and DAPI (blue, nuclear).



IHC staining of FFPE human skin with MICA antibody. HIER: boil tissue sections in pH6, 10mM citrate buffer, for 10-20 min and allow to cool before testing.

Description

MICA is the highly polymorphic MHC (HLA) class I chain-related gene A. The protein product is expressed on the cell surface, although unlike canonical class I molecules does not seem to associate with beta-2-microglobulin. It is thought that MICA functions as a stress-induced antigen that is broadly recognized by intestinal epithelial gamma delta T cells.

Application Notes

Titration of the MICA antibody may be required due to differences in protocols and secondary/substrate sensitivity.

Immunogen

A portion of amino acids 68-97 from the human protein was used as the immunogen for this MICA antibody.

Storage

Aliquot the MICA antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.