


MNAT1 Antibody (R32131)


Catalog No.	Formulation	Size
R32131	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug

Bulk quote request


Availability	1-3 business days
Species Reactivity	Human, Mouse, Rat
Format	Antigen affinity purified
Host	Rabbit
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit IgG
Purity	Antigen affinity
Buffer	Lyophilized from 1X PBS with 2.5% BSA and 0.025% sodium azide
UniProt	P51948
Localization	Nuclear
Applications	Western Blot : 0.5-1ug/ml Immunohistochemistry (FFPE) : 1-2ug/ml Immunofluorescence (FFPE) : 5ug/ml Flow Cytometry : 1-3ug/million cells
Limitations	This MNAT1 antibody is available for research use only.




Immunofluorescent staining of FFPE human A431 cells with MNAT1 antibody (green) and DAPI nuclear stain (blue). HIER: steam section in pH6 citrate buffer for 20 min.




IHC testing of FFPE human testis with MNAT1 antibody. HIER: Boil the paraffin sections in pH 6, 10mM citrate buffer for 20 minutes and allow to cool prior to staining.




IHC testing of FFPE mouse testis with MNAT1 antibody. HIER: Boil the paraffin sections in pH 6, 10mM citrate buffer for 20 minutes and allow to cool prior to staining.



IHC testing of FFPE rat testis with MNAT1 antibody. HIER: Boil the paraffin sections in pH 6, 10mM citrate buffer for 20 minutes and allow to cool prior to staining.



Western blot testing of 1) human HeLa, 2) human A431, 3) human A375, 4) human MCF7, 5) rat testis, 6) rat L6, 7) mouse testis and 8) mouse NIH 3T3 cell lysate with MNAT1 antibody. Expected molecular weight ~36 kDa.



Flow cytometry testing of human A431 cells with MNAT1 antibody at 1ug/million cells (blocked with goat sera); Red=cells alone, Green=isotype control, Blue= MNAT1 antibody.

Description

CDK-activating kinase assembly factor MAT1 is an enzyme that in humans is encoded by the MNAT1 gene. The protein encoded by this gene, along with cyclin H and CDK7, forms the CDK-activating kinase (CAK) enzymatic complex. This complex activates several cyclin-associated kinases and can also associate with TFIIF to activate transcription by RNA polymerase II. Two transcript variants encoding different isoforms have been found for this gene.

Application Notes

Optimal dilution of the MNAT1 antibody should be determined by the researcher.

Immunogen

Amino acids 92-278 of human MNAT1 were used as the immunogen for the MNAT1 antibody.

Storage

After reconstitution, the MNAT1 antibody can be stored for up to one month at 4oC. For long-term, aliquot and store at -20oC. Avoid repeated freezing and thawing.