


## MOBKL3 Antibody / PREI3 (R34918)


| Catalog No.  | Formulation   | Size   |
|--------------|---|--------|
| R34918-100UG | 0.5 mg/ml in 1X TBS, pH7.3, with 0.5% BSA (US sourced) and 0.02% sodium azide | 100 ug |

[Bulk quote request](#)


|                             |  |
|-----------------------------|--|
| <b>Availability</b>         | 1-3 business days  |
| <b>Species Reactivity</b>   | Human, Mouse, Rat  |
| <b>Predicted Reactivity</b> | Dog, Cow   |
| <b>Format</b>               | Antigen affinity purified  |
| <b>Host</b>                 | Goat   |
| <b>Clonality</b>            | Polyclonal (goat origin)   |
| <b>Isotype</b>              | Goat Ig  |
| <b>Purity</b>               | Antigen affinity   |
| <b>Gene ID</b>              | 25843  |
| <b>Applications</b>         | Western Blot : 0.3-1ug/ml<br>Immunohistochemistry (FFPE) : 5-10ug/ml<br>ELISA (peptide) LOD : 1:4000 |
| <b>Limitations</b>          | This MOBKL3 antibody is available for research use only.   |



Western blot testing of human frontal cortex lysate with MOBKL3 antibody at 0.3ug/ml.  
Predicted/observed molecular weight: ~26 kDa.



IHC staining of FFPE human cerebral cortex with MOBKL3 antibody at 5ug/ml. HIER: steamed with pH6 citrate buffer, AP-staining.



Western blot testing with MOBKL3 antibody: Lane A: mouse brain at 0.3ug/ml; Lane B: rat brain at 1ug/ml. Predicted/observed molecular weight: ~26 kDa.

## Description

Additional name(s) for this target protein: Mps One Binder kinase activator-like 3, MOB1; PREI3; Preimplantation protein 3; MOB3; MOB4

## Application Notes

Optimal dilution of the MOBKL3 antibody should be determined by the researcher.

## Immunogen

Amino acids NSNKYFPSRVSİKES were used as the immunogen for this MOBKL3 antibody.

## Storage

Aliquot and store the MOBKL3 antibody at -20oC.