


MRP8 Antibody / Multidrug resistance-associated protein 8 / ABCC11 [clone ABCC11/2438] (V5611)


Catalog No.	Formulation	Size
V5611-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	100 ug
V5611-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	20 ug
V5611SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

Bulk quote request

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Host	Mouse
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG2b, kappa
Clone Name	ABCC11/2438
Purity	Protein A/G affinity
UniProt	Q96J66
Localization	Membrane
Applications	Immunohistochemistry (FFPE) : 1-2ug/ml
Limitations	This MRP8 antibody is available for research use only.



IHC staining of FFPE triple negative breast cancer tissue with MRP8 antibody (clone ABCC11/2438). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



SDS-PAGE analysis of purified, BSA-free MRP8 antibody (clone ABCC11/2438) as confirmation of integrity and purity.

Description

ATP-binding cassette (ABC) transporters belong to an evolutionarily conserved family of proteins that catalyze the transport of molecules across extra- and intracellular membranes through the energy of ATP hydrolysis. ABC genes comprise seven subfamilies, designated ABC1, MDR/TAP, MRP, ALD, OABP, GCN20 and White. The complete human ABCC subfamily has 12 identified members (ABCC1-12), nine from the multidrug resistance-like subgroup, two from the sulfonylurea receptor subgroup, and the CFTR gene. The human ABCC11 gene maps to chromosome 16q12.1 and encodes a 1,382 amino acid protein. The human ABCC12 gene maps to chromosome 16q12.1 and encodes a 1,359 amino acid protein. Transcripts of ABCC11 and ABCC12 genes are present in various adult human tissues, including liver, lung and kidney, and also in several fetal tissues. Their chromosomal localization, potential function and expression patterns identify them as candidates for paroxysmal kinesigenic choreoathetosis, a disorder characterized by attacks of involuntary movements and postures, chorea and dystonia. Other inherited disorders where ABC transporters are implicated include cystic fibrosis, neurological disease, retinal degeneration, cholesterol and bile transport defects, anemia and drug response.

Application Notes

Optimal dilution of the MRP8 antibody should be determined by the researcher.

Immunogen

A recombinant fragment (within amino acids 400-600) of human ABCC11/MRP8 protein was used as the immunogen for the MRP8 antibody.

Storage

Aliquot the MRP8 antibody and store frozen at -20°C or colder. Avoid repeated freeze-thaw cycles.