


MSLN Antibody / Mesothelin [clone MSLN/3387] (V8693)


Catalog No.	Formulation	Size
V8693-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V8693-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V8693SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

Bulk quote request


Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG1, kappa
Clone Name	MSLN/3387
Purity	Protein G affinity chromatography
UniProt	Q13421
Localization	Cytoplasmic, cell surface, secreted
Applications	Immunohistochemistry (FFPE) : 1-2ug/ml for 30 minutes at RT Western Blot : 2-4ug/ml
Limitations	This MSLN antibody is available for research use only.




IHC staining of FFPE human mesothelioma with MSLN antibody. HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



IHC staining of FFPE human mesothelioma with MSLN antibody. HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



Analysis of HuProt(TM) microarray containing more than 19,000 full-length human proteins using MSLN antibody. These results demonstrate the foremost specificity of the MSLN/3387 mAb. Z- and S- score: The Z-score represents the strength of a signal that an antibody (in combination with a fluorescently-tagged anti-IgG secondary Ab) produces when binding to a particular protein on the HuProt(TM) array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If the targets on the HuProt(TM) are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-scores. The S-score therefore represents the relative target specificity of an Ab to its intended target.



Western blot testing of human OVCAR-3 cell lysate with MSLN antibody. Predicted molecular weight ~70 kDa (precursor), ~40 kDa (processed form).

Description

Mesothelin is a 40kDa glycosyl-phosphatidylinositol-anchored glycoprotein cleaved from a 69kDa precursor protein. Mesothelin immunoreactivity is high in cancers of the ovary (serous papillary, endometrioid and undifferentiated) and pancreas, with less frequent staining seen in adenocarcinomas of the endometrium, lung and stomach/esophagus. Mesothelin is one of the most sensitive markers for mesothelioma.

Application Notes

Optimal dilution of the MSLN antibody should be determined by the researcher.

Immunogen

A portion of amino acids 273-407 from the human protein was used as the immunogen for the MSLN antibody.

Storage

Store the MSLN antibody at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).