


## MUC16 Antibody / CA125 [clone CA125/8893] (V6040)


Catalog No.	Formulation	Size
V6040-100UG	0.2 mg/ml in 1X PBS with 0.05% BSA, 0.05% sodium azide	100 ug
V6040-20UG	0.2 mg/ml in 1X PBS with 0.05% BSA, 0.05% sodium azide	20 ug
V6040SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

[Bulk quote request](#)


<b>Species Reactivity</b>	Human
<b>Format</b>	Purified
<b>Host</b>	Mouse
<b>Clonality</b>	Monoclonal (mouse origin)
<b>Isotype</b>	Mouse IgG2b, kappa
<b>Clone Name</b>	CA125/8893
<b>UniProt</b>	Q8WXI7
<b>Localization</b>	Cell membrane, Extracellular space, Secreted
<b>Applications</b>	Immunohistochemistry (FFPE) : 1-2ug/ml
<b>Limitations</b>	This MUC16/CA125 antibody is available for research use only.



Formalin-fixed, paraffin-embedded human colon carcinoma stained with MUC16 Mouse Monoclonal Antibody (CA125/8893). Inset: PBS instead of primary antibody; secondary only negative control.



SDS-PAGE Analysis of Purified MUC16 Mouse Monoclonal Antibody (CA125/8893). Confirmation of Purity and Integrity of Antibody.



Formalin-fixed, paraffin-embedded human ovarian carcinoma stained with MUC16 Mouse Monoclonal Antibody (CA125/8893). Inset: PBS instead of primary antibody; secondary only negative control.

## Application Notes

Optimal dilution of the MUC16/CA125 antibody should be determined by the researcher.

## Immunogen

A recombinant fragment (around amino acids 1700-2000) of human MUC16 protein (exact sequence is proprietary) was used as the immunogen for the MUC16/CA125 antibody.

## Storage

MUC16/CA125 antibody with sodium azide - store at 2 to 8°C; antibody without sodium azide - store at -20 to -80°C.