


## MUC16 Antibody / Ovarian cancer antigen 125 [clone CA125/13895R] (V6042)

Catalog No.	Formulation	Size
V6042-100UG	0.2 mg/ml in 1X PBS with 0.05% BSA, 0.05% sodium azide	100 ug
V6042-20UG	0.2 mg/ml in 1X PBS with 0.05% BSA, 0.05% sodium azide	20 ug
V6042SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

Recombinant **RABBIT MONOCLONAL**


[Bulk quote request](#)

<b>Species Reactivity</b>	Human
<b>Format</b>	Purified
<b>Host</b>	Rabbit
<b>Clonality</b>	Recombinant Rabbit Monoclonal
<b>Isotype</b>	Rabbit IgG, kappa
<b>Clone Name</b>	CA125/13895R
<b>Purity</b>	Protein A affinity
<b>UniProt</b>	Q8WXI7
<b>Localization</b>	Cell membrane, Extracellular space, Secreted
<b>Applications</b>	Immunohistochemistry (FFPE) : 1-2ug/ml
<b>Limitations</b>	This MUC16/Ovarian cancer antigen 125 antibody is available for research use only.



Formalin-fixed, paraffin-embedded human ovarian carcinoma stained with Mucin 16 / CA125 Recombinant Rabbit Monoclonal Antibody (CA125/13895R). Inset: PBS instead of primary antibody; secondary only negative control.

Purity: SDS-PAGE



SDS-PAGE under non-reducing (NR) and reducing (R) conditions. The gel was stained with Coomassie Blue. The purity of the protein is 96%.

SDS-PAGE Analysis of Purified Mucin 16 / CA125 Recombinant Rabbit Monoclonal Antibody (CA125/13895R). Confirmation of Purity and Integrity of Antibody.

## Application Notes

1. Optimal dilution of the MUC16/Ovarian cancer antigen 125 antibody should be determined by the researcher.
2. This MUC16/Ovarian cancer antigen 125 antibody is recombinantly produced by expression in CHO cells.

## Immunogen

Recombinant human MUC16 protein was used as the immunogen for the MUC16/Ovarian cancer antigen 125 antibody.

## Storage

MUC16/Ovarian cancer antigen 125 antibody with sodium azide - store at 2 to 8°C; antibody without sodium azide - store at -20 to -80°C.