


MUC3 Antibody [clone MUCN3-1] (V7118)


Catalog No.	Formulation	Size
V7118-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V7118-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V7118SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug
V7118IHC-7ML	Prediluted in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide; *For IHC use only*	7 ml

Bulk quote request

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG2a, kappa
Clone Name	MUCN3-1
Purity	Protein G affinity chromatography
UniProt	Q02505
Localization	Cytoplasmic and cell surface
Applications	Immunohistochemistry (FFPE) : 1-2ug/ml for 30 min at RT (1) Prediluted IHC Only Format : incubate for 30 min at RT (2)
Limitations	This MUC3 antibody is available for research use only.



IHC testing of FFPE human gastric carcinoma with MUC3 antibody (clone MUCN3-1). Staining of formalin-fixed tissues requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 min.



IHC testing of FFPE human colon carcinoma with MUC3 antibody (clone MUCN3-1). Staining of formalin-fixed tissues requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 min.

Description

The Mucins are a family of highly glycosylated, secreted proteins with a basic structure consisting of a variable number of tandem repeats. Mucin-3/MUC3 is a major glycoprotein component of a variety of mucus gels. Thought to provide a protective, lubricating barrier against particles and infectious agents at mucosal surfaces. May be involved in ligand binding and intracellular signaling. [UniProt]

Application Notes

Optimal dilution of the MUC3 antibody should be determined by the researcher.

1. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.

Immunogen

Amino acids (C-HSTPSFTSSITTTETTTSHSTPSFTSSITTTETTS) which contains two of the MUC3 tandem repeats was used as the immunogen for the MUC3 antibody.

Storage

Store the MUC3 antibody at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).