


MUC5AC Antibody [clone 58M1] (V2734)

Catalog No.	Formulation	Size
V2734-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V2734-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V2734SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

Citations ((4)
-------------	-----

Bulk quote request

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG1, kappa
Clone Name	58M1
Purity	Protein G affinity chromatography
UniProt	P98088
Localization	Cytoplasmic
Applications	ELISA : 1-5ug/ml for coating (order BSA/sodium azide-free format)
Limitations	This MUC5AC antibody is available for research use only.



SDS-PAGE analysis of purified, BSA-free MUC5AC antibody (clone 58M1) as confirmation of integrity and purity.

Description

This mAb recognizes the peptide core of gastric mucin M1 (recently identified as Mucin 5AC). Its epitope is destroyed by beta-mercaptoethanol but not by periodate treatment. This mucin is present in primary ovarian mucinous cancer but

usually absent in colorectal adenocarcinoma, thus showing an expression pattern opposite to MUC2. Together with a panel of antibodies, Anti-MUC5AC may be useful for differential identification of primary mucinous ovarian tumors from colon adenocarcinoma metastatic to the ovary. MUC5AC antibodies may also be useful for identification of intestinal metaplasia as well as in the identification of pancreatic carcinoma and pre-cancerous changes vs. normal pancreas.

Application Notes

Optimal dilution of the MUC5AC antibody should be determined by the researcher.

Immunogen

An M1 mucin preparation from the fluid of an ovarian mucinous cyst belonging to an O Le(a-b) patient was used as the immunogen for the MUC5AC antibody.

Storage

Store the MUC5AC antibody at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).