


## MyD88 Antibody (F55059)


Catalog No.	Formulation	Size
F55059-0.2ML	In 1X PBS, pH 7.4, with 0.09% sodium azide	0.2 ml
F55059-0.05ML	In 1X PBS, pH 7.4, with 0.09% sodium azide	0.05 ml

[Bulk quote request](#)


<b>Availability</b>	1-3 business days
<b>Species Reactivity</b>	Human
<b>Format</b>	Antigen affinity purified
<b>Clonality</b>	Polyclonal (rabbit origin)
<b>Isotype</b>	Rabbit Ig
<b>Purity</b>	Antigen affinity
<b>UniProt</b>	Q99836
<b>Applications</b>	Western Blot : 1:1000-1:2000 Immunohistochemistry (FFPE) : 1:10-1:50 Flow Cytometry : 1:10-1:50 per million cells in 0.1ml
<b>Limitations</b>	This MyD88 antibody is available for research use only.



MyD88 antibody flow cytometric analysis of fixed and permeabilized human K562 cells (green) compared to a  $\lambda$  negative control (blue). FITC-conjugated goat-anti-rabbit secondary Ab was used for the analysis.



Western blot testing of human HT-29 cell lysate with MyD88 antibody. Predicted molecular weight ~33 kDa.



IHC analysis of FFPE human lung carcinoma stained with MyD88 antibody. HIER: steam section in pH6 citrate buffer for 20 min and allow to cool prior to staining.

## Description

Myeloid differentiation primary response protein MyD88 is an adapter protein involved in the Toll-like receptor and IL-1 receptor signaling pathway in the innate immune response. It acts via IRAK1, IRAK2 and TRAF6, leading to NF-kappa-B activation, cytokine secretion and the inflammatory response and increases IL-8 transcription. It may be involved in myeloid differentiation.

## Application Notes

Titration of the MyD88 antibody may be required due to differences in protocols and secondary/substrate sensitivity.

## Immunogen

A portion of amino acids 136-164 from the human protein was used as the immunogen for this MyD88 antibody.

## Storage

Aliquot the MyD88 antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.