

Myosin 7 Antibody / MYH7 [clone MYH7/9184] (V5475)


Catalog No.	Formulation	Size
V5475-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	100 ug
V5475-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	20 ug
V5475SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

Bulk quote request

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Host	Mouse
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG2b, kappa
Clone Name	MYH7/9184
Purity	Protein A/G affinity
UniProt	P12883
Localization	Cytoplasm
Applications	Immunohistochemistry (FFPE) : 1-2ug/ml
Limitations	This Myosin 7 antibody is available for research use only.



IHC staining of FFPE human skeletal muscle tissue with Myosin 7 antibody (clone MYH7/9184). Inset: PBS used in place of primary Ab (secondary Ab negative control). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



SDS-PAGE analysis of purified, BSA-free Myosin 7 antibody (clone MYH7/9184) as confirmation of integrity and purity.

Description

Myosin 7 antibody is used to detect Myosin 7/MYH7. Myosin heavy chains are ubiquitous Actin-based motor proteins that convert the chemical energy derived from ATP hydrolysis into the mechanical energy that drives diverse motile processes in eukaryotic cells, including cytokinesis, vesicular transport and cellular locomotion. Muscle myosin is a heterohexamer consisting of two myosin heavy chains and two associated nonidentical pairs of myosin light chains. The seven myosin heavy chain isoforms that predominate in mammalian skeletal muscles include two developmental isoforms, MHC-embryonic (MYH3) and MHC-perinatal (MYH8); three adult skeletal muscle isoforms, MHC IIa (MYH2), MHC IIb (MYH4) and MHC IIx/d (MYH1); and MHC-?/slow (MYH7 or MHC-?), which is also expressed in cardiac muscle. Research indicates that mutations of the MYH7 gene causes hypertrophic cardiomyopathy.

Application Notes

Optimal dilution of the Myosin 7 antibody should be determined by the researcher.

Immunogen

A recombinant fragment (within amino acids 1150-1350) of human Myosin 7 protein was used as the immunogen for the Myosin 7 antibody.

Storage

Aliquot the Myosin 7 antibody and store frozen at -20°C or colder. Avoid repeated freeze-thaw cycles.