


## MYRIP Antibody (R34377)


Catalog No.	Formulation	Size
R34377-100UG	0.5 mg/ml in 1X TBS, pH7.3, with 0.5% BSA (US sourced) and 0.02% sodium azide	100 ug

[Bulk quote request](#)


<b>Availability</b>	1-3 business days
<b>Species Reactivity</b>	Human
<b>Format</b>	Antigen affinity purified
<b>Host</b>	Goat
<b>Clonality</b>	Polyclonal (goat origin)
<b>Isotype</b>	Goat Ig
<b>Purity</b>	Antigen affinity
<b>Gene ID</b>	25924
<b>Localization</b>	Cytoplasmic
<b>Applications</b>	Western Blot : 0.3-1ug/ml IHC (FFPE) : 3.75ug/ml ELISA (peptide) LOD : 1:64000
<b>Limitations</b>	This MYRIP antibody is available for research use only.




Western blot testing of human amygdala lysate with MYRIP antibody at 0.3ug/ml.  
Predicted molecular weight: ~95 kDa.



IHC testing of FFPE human cortex (brain) tissue with MYRIP antibody at 3.75ug/ml. Required HIER: steamed antigen retrieval with pH6 citrate buffer; AP-staining.



IHC testing of FFPE human colon tissue with MYRIP antibody at 3.75ug/ml. Required HIER: steamed antigen retrieval with pH6 citrate buffer; AP-staining.



IHC testing of FFPE human breast tissue with MYRIP antibody at 3.75ug/ml. Required HIER: steamed antigen retrieval with pH6 citrate buffer; AP-staining.

## Description

Additional name(s) for this target protein: Myosin VIIA and Rab interacting protein, SLAC2C

## Application Notes

Optimal dilution of the MYRIP antibody should be determined by the researcher.

## Immunogen

Amino acids KDLMEPALESAVMY were used as the immunogen for this MYRIP antibody.

## Storage

Aliquot and store the MYRIP antibody at -20°C.