


## Myxovirus resistance protein 1 Antibody / MX1 / MxA [clone MX1/7530] (V4845)


Catalog No.	Formulation	Size
V4845-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	100 ug
V4845-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	20 ug
V4845SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

### Bulk quote request


Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Host	Mouse
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG2, kappa
Clone Name	MX1/7530
Purity	Protein A/G affinity
UniProt	P20591
Localization	Cytoplasm
Applications	Immunohistochemistry (FFPE) : 1-2ug/ml for 30 min at RT
Limitations	This Myxovirus resistance protein 1 antibody is available for research use only.



IHC staining of FFPE human kidney tissue with Myxovirus resistance protein 1 antibody (clone MX1/7530). Inset: PBS used in place of primary Ab (secondary Ab negative control). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



IHC staining of FFPE human ovarian cancer tissue with Myxovirus resistance protein 1 antibody (clone MX1/7530). Inset: PBS used in place of primary Ab (secondary Ab negative control). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



SDS-PAGE analysis of purified, BSA-free Myxovirus resistance protein 1 antibody (clone MX1/7530) as confirmation of integrity and purity.

## Description

The Dynamin family of microtubule-associated proteins function as GTPases that are involved in microtubule bundling and endocytosis. In mice, Mx2 (myxovirus resistance protein 2) and Mx1 (myxovirus resistance protein 1) are members of the Dynamin family that are involved in the immune response to viral infections. Localized to the cytoplasm, Mx2 contains one GED domain and is expressed in response to viral infection or treatment by IFN- $\alpha$ /IFN- $\beta$ . Once expression is induced, Mx2 accumulates in the cytoplasm and inhibits the replication of vesicular stomatitis virus (VSV), thereby conferring resistance to VSV infection. Unlike Mx2, Mx1 is localized to the nucleus where, upon induction by IFN- $\alpha$ /IFN- $\beta$ , it provides selective resistance to infection by the highly lethal H5N1 influenza virus. In humans, MxA and MxB function in a similar manner to Mx1 and Mx2, conferring resistance to specific target viruses. Mx3 is a rat-specific member of the myxovirus resistance protein family.

## Application Notes

Optimal dilution of the Myxovirus resistance protein 1 antibody should be determined by the researcher.

## Immunogen

A recombinant partial protein sequence (within amino acids 400-592) from the human protein was used as the immunogen for the Myxovirus resistance protein 1 antibody.

## Storage

Aliquot the Myxovirus resistance protein 1 antibody and store frozen at -20°C or colder. Avoid repeated freeze-thaw cycles.