


## N4BP1 Antibody / NEDD4-binding protein 1 / KIAA0615 (RQ8184)


| Catalog No. | Formulation   | Size   |
|-------------|---|--------|
| RQ8184      | 0.5mg/ml if reconstituted with 0.2ml sterile DI water | 100 ug |

### Bulk quote request


|                           |   |
|---------------------------|---|
| <b>Availability</b>       | 1-3 business days   |
| <b>Species Reactivity</b> | Human   |
| <b>Format</b>             | Antigen affinity purified   |
| <b>Host</b>               | Rabbit  |
| <b>Clonality</b>          | Polyclonal (rabbit origin)  |
| <b>Isotype</b>            | Rabbit IgG  |
| <b>Purity</b>             | Antigen affinity purified   |
| <b>Buffer</b>             | Lyophilized from 1X PBS with 2% Trehalose   |
| <b>UniProt</b>            | O75113  |
| <b>Localization</b>       | Nuclear, cytoplasmic  |
| <b>Applications</b>       | Western Blot : 0.5-1ug/ml<br>Immunofluorescence : 5ug/ml<br>Flow Cytometry : 1-3ug/million cells<br>Direct ELISA : 0.1-0.5ug/ml |
| <b>Limitations</b>        | This N4BP1 antibody is available for research use only.   |



Immunofluorescent staining of FFPE human HeLa cells with N4BP1 antibody (red) and DAPI nuclear stain (blue). HIER: steam section in pH6 citrate buffer for 20 min.



Western blot testing of human 1) HepG2, 2) RT4 and 3) Jurkat cell lysate with N4BP1 antibody. Predicted molecular weight ~100 kDa.



Flow cytometry testing of fixed and permeabilized human JK cells with N4BP1 antibody at 1ug/million cells (blocked with goat sera); Red=cells alone, Green=isotype control, Blue= N4BP1 antibody.

## Description

Nedd4-binding partner-1 (N4BP1) has been identified as a protein interactor and a substrate of the homologous to E6AP C terminus (HECT) domain-containing E3 ubiquitin-protein ligase (E3), Nedd4. Here, we describe a previously unrecognized functional interaction between N4BP1 and Itch, a Nedd4 structurally related E3, which contains four WW domains, conferring substrate-binding activity. We show that N4BP1 association with the second WW domain (WW2) of Itch interferes with E3 binding to its substrates. In particular, we found that N4BP1 and p73<sup>Δ</sup>, a target of Itch-mediated ubiquitin/proteasome proteolysis, share the same binding site. By competing with p73<sup>Δ</sup> for binding to the WW2 domain, N4BP1 reduces the ability of Itch to recruit and ubiquitylate p73<sup>Δ</sup> and inhibits Itch autoubiquitylation activity both in *in vitro* and *in vivo* ubiquitylation assays.

## Application Notes

Optimal dilution of the N4BP1 antibody should be determined by the researcher.

## Immunogen

*E. coli*-derived recombinant human protein (amino acids M1-K885) was used as the immunogen for the N4BP1 antibody.

## Storage

After reconstitution, the N4BP1 antibody can be stored for up to one month at 4°C. For long-term, aliquot and store at -20°C. Avoid repeated freezing and thawing.