


# Napsin A Antibody [clone NAPSA/3308] (V7604BTN)


Catalog No.	Formulation	Size
V7604BTN	0.1 mg/ml with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	500 ul

## **Bulk quote request**


Availability	1-3 business days
Species Reactivity	Human
Format	Biotin Conjugate
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG1, kappa
Clone Name	NAPSA/3308
Purity	Protein G affinity chromatography
UniProt	O96009
Localization	Cytoplasmic
Applications	Immunohistochemistry (FFPE): 2-4ug/ml for 30 minutes at RT
Limitations	This Napsin A antibody is available for research use only.




IHC staining of FFPE human lung adenocarcinoma with biotin-conjugated Napsin A antibody (clone NAPSA/3308). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



IHC staining of FFPE human lung adenocarcinoma with biotin-conjugated Napsin A antibody (clone NAPSA/3308). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



Analysis of HuProt(TM) microarray containing more than 19,000 full-length human proteins using Napsin A antibody (clone NAPSA/3308). These results demonstrate the foremost specificity of the NAPSA/3308 mAb. Z- and S- score: The Z-score represents the strength of a signal that an antibody (in combination with a fluorescently-tagged antilgG secondary Ab) produces when binding to a particular protein on the HuProt(TM) array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If the targets on the HuProt(TM) are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-scores. The S-score therefore represents the relative target specificity of an Ab to its intended target.



SDS-PAGE analysis of purified, BSA-free Napsin A antibody (clone NAPSA/3308) as confirmation of integrity and purity.

### **Description**

Napsin is a pepsin-like aspartic proteinase connected with maturation of surfactant protein B. There are two closely related napsins, napsin A and napsin B. Napsin A is expressed as a single chain protein. Immunohistochemical studies revealed high expression levels of napsin A in human lung and kidney but low expression in spleen. Napsin A is expressed in type II pneumocytes and in adenocarcinomas of lung. The high specificity expression of napsin A in adenocarcinomas of lung is useful to distinguish primary lung adenocarcinomas from adenocarcinomas of other organs.

## **Application Notes**

Optimal dilution of the Napsin A antibody should be determined by the researcher.

#### **Immunogen**

Amino acids 189-299 from the human protein were used as the immunogen for the Napsin A antibody.

#### **Storage**

Store the Napsin A antibody at 2-8oC (up to one month) or aliquot and store at -20oC (longer term).