
NCAM Antibody NCAM1/784 / Mouse Monoclonal [clone NCAM1/784] (V2753)

Catalog No.	Formulation	Size
V2753-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V2753-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V2753SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug
V2753IHC-7ML	Prediluted in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide; *For IHC use only*	7 ml

Bulk quote request

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Host	Mouse
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG1, kappa
Clone Name	NCAM1/784
Purity	Protein G affinity chromatography
UniProt	P13591
Localization	Cell surface, cytoplasmic
Applications	Flow cytometry: 0.5-1ug/10^6 cells Immunofluorescence: 1-2ug/ml Immunohistochemistry (FFPE): 0.5-1ug/ml for 30 min at RT
Limitations	This NCAM Antibody NCAM1/784 is available for research use only.



IHC: Formalin-fixed, paraffin-embedded human colon stained with NCAM Antibody NCAM1/784.

Description

NCAM Antibody NCAM1/784 reacts with an extracellular domain (close to transmembrane) of CD56/NCAM. Three isoforms of neural cell adhesion molecule (NCAM) are produced by differential splicing of the RNA transcript from a single gene. The 135kDa isoform is the basic molecule, which is glycosylated or sialylated to produce the mature species. Anti-CD56 recognizes two proteins of the neural cell adhesion molecule, the basic molecule expressed on most neuroectodermally derived tissues and neoplasms (e.g. retinoblastoma, medulloblastomas, astrocytomas, neuroblastomas, and small cell carcinomas). It is also expressed on some mesodermally derived tumors (rhabdomyosarcoma). Anti-CD56 plays an important role in the diagnosis of nodal and nasal NK/T-cell lymphomas.

For additional information on NCAM1/CD56 and its roles in neural cell adhesion, neuronal development, and cell signaling, see our NCAM1 Antibody / Neural Cell Adhesion Molecule 1 Antibody page.

Application Notes

Optimal dilution of the NCAM Antibody NCAM1/784 should be determined by the researcher.

1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min
2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.

Immunogen

Recombinant human protein was used as the immunogen for the NCAM antibody.

Storage

Store the NCAM antibody at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).

Copyright © NSJ Bioreagents. All rights reserved.