# NCAM Antibody [clone rNCAM1/8655] (V4202)

| Catalog No. | Formulation | Size |
|---|---|---|
| V4202-100UG | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide | 100 ug |
| V4202-20UG | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide | 20 ug |
| V4202SAF-100UG | 1 mg/ml in 1X PBS; BSA free, sodium azide free | 100 ug |

Recombinant | MOUSE MONOCLONAL

**Bulk quote request**

| | |
|---|---|
| **Availability** | 1-3 business days |
| **Species Reactivity** | Human |
| **Format** | Purified |
| **Host** | Mouse |
| **Clonality** | Recombinant Mouse Monoclonal |
| **Isotype** | Mouse IgG1, kappa |
| **Clone Name** | rNCAM1/8655 |
| **Purity** | Protein A/G affinity |
| **UniProt** | P13591, P13592 |
| **Localization** | Cell surface |
| **Applications** | Immunohistochemistry (FFPE) : 1-2ug/ml for 30 minutes at RT |
| **Limitations** | This NCAM antibody is available for research use only. |



IHC staining of FFPE human colon tissue with CD56 antibody (clone rNCAM1/8655). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.

## Description

This mAb reacts with an extracellular domain (close to transmembrane) of CD56/NCAM. Three isoforms of neural cell adhesion molecule (NCAM) are produced by differential splicing of the RNA transcript from a single gene. The 135kDa

isoform is the basic molecule, which is glycosylated or sialylated to produce the mature species. Anti-CD56 recognizes two proteins of the neural cell adhesion molecule, the basic molecule expressed on most neuroectodermally derived tissues and neoplasms (e.g. retinoblastoma, medulloblastomas, astrocytomas, neuroblastomas, and small cell carcinomas). It is also expressed on some mesodermally derived tumors (rhabdomyosarcoma). Anti-CD56 plays an important role in the diagnosis of nodal and nasal NK/T-cell lymphomas.

## Application Notes

Optimal dilution of the NCAM antibody should be determined by the researcher.

## Immunogen

Recombinant human NCAM1 protein was used as the immunogen for the NCAM antibody.

## Storage

Aliquot the NCAM antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.