

Neurofilament Antibody Cocktail [clone RT-97 + NR-4] (V3088)


Catalog No.	Formulation	Size
V3088-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V3088-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V3088SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug
V3088IHC-7ML	Prediluted in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide; *For IHC use only*	7 ml




Citations (4)

[Bulk quote request](#)


Availability	1-3 business days
Species Reactivity	Human, Mouse, Rat
Format	Purified
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG1, kappa
Clone Name	RT-97 + NR-4
Purity	Protein G affinity chromatography
UniProt	P12036
Localization	Cytoplasmic, membranous
Applications	Flow Cytometry : 0.5-1ug/10 ⁶ cells Immunohistochemistry (FFPE) : 0.25-0.5ug/ml for 30 min at RT
Limitations	This Neurofilament antibody cocktail is available for research use only.



IHC: Formalin-fixed, paraffin-embedded human cerebellum stained with Neurofilament antibody (RT-97 + NR-4).



IHC: Formalin-fixed, paraffin-embedded rat cerebellum stained with Neurofilament antibody (RT-97 + NR-4).



Flow cytometry testing of permeabilized human HEK293 cells with Neurofilament antibody cocktail (clone RT-97 + NR-4); Red=isotype control, Blue= Neurofilament antibody cocktail.

Description

This mAb reacts with a 200kDa and 68kDa protein, identified as heavy and light sub-units of neurofilaments (NF-H & NF-L). Neurofilaments make up the main structural elements of axons and dendrites and are found in neurons, peripheral nerves, and sympathetic ganglion cells. Neurofilaments consist of three major subunits with molecular weights of 68kDa (NF-L), 160kDa (NF-M) and 200kDa (NF-H). Anti-neurofilament stains a number of neural, neuroendocrine, and endocrine tumors. Neuromas, ganglioneuromas, gangliogliomas, ganglioneuroblastomas, and neuroblastomas stain positively for anti-neurofilament. Neurofilaments are also present in paragangliomas as well as adrenal and extra-adrenal pheochromocytomas. Carcinoids, neuroendocrine carcinomas of the skin, and cell carcinomas of the lung also express neurofilament.

Application Notes

Optimal dilution of the Neurofilament antibody cocktail should be determined by the researcher.

1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min.
2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.

Immunogen

Triton-X 100 insoluble proteins of rat brain (RT-97) and Neurofilaments from porcine spinal cord (NR-4) were used as the immunogen for the Neurofilament antibody cocktail.

Storage

Store the Neurofilament antibody cocktail at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).

