


NFATC2 Antibody (R35408)


Catalog No.	Formulation	Size
R35408-100UG	0.5 mg/ml in 1X TBS, pH7.3, with 0.5% BSA (US sourced) and 0.02% sodium azide	100 ug

[Bulk quote request](#)

Availability	1-3 business days
Species Reactivity	Human
Predicted Reactivity	Mouse, Rat, Dog, Cow
Format	Antigen affinity purified
Host	Goat
Clonality	Polyclonal (goat origin)
Isotype	Goat Ig
Purity	Antigen affinity
Gene ID	4773
Localization	Cytoplasmic
Applications	Western Blot : 0.3-1ug/ml Immunohistochemistry (FFPE) : 3-6ug/ml ELISA (peptide) LOD : 1:64000
Limitations	This NFATC2 antibody is available for research use only.



Western blot testing of MOLT4 lysate with NFATC2 antibody at 0.3ug/ml. Predicted molecular weight: ~100 kDa.



IHC staining of FFPE human spleen with NFATC2 antibody at 4ug/ml. HIER: steamed with pH6 citrate buffer, AP-staining.

Description

Additional name(s) for this target protein: Nuclear factor of activated T-cells, cytoplasmic, calcineurin-dependent 2

Application Notes

Optimal dilution of the NFATC2 antibody should be determined by the researcher.

Immunogen

Amino acids QQFRTGLSSPDARYQ were used as the immunogen for this NFATC2 antibody.

Storage

Aliquot and store the NFATC2 antibody at -20oC.