


NKX2.1 Antibody / TFF-1 / Thyroid transcription factor 1 [clone NX2.1/9033] (V5801)


Catalog No.	Formulation	Size
V5801-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	100 ug
V5801-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	20 ug
V5801SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

Bulk quote request


Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Host	Mouse
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG2a, kappa
Clone Name	NX2.1/9033
Purity	Protein G affinity
UniProt	P43699
Localization	Nucleus
Applications	Immunohistochemistry (FFPE) : 1-2ug/ml
Limitations	This NKX2.1 antibody is available for research use only.



IHC staining of FFPE human thyroid tissue with NKX2.1 antibody (clone NX2.1/9033). Inset: PBS used in place of primary Ab (secondary Ab negative control). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



IHC staining of FFPE human thyroid tissue with NKX2.1 antibody (clone NX2.1/9033).
HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



SDS-PAGE analysis of purified, BSA-free NKX2.1 antibody (clone NX2.1/9033) as confirmation of integrity and purity.

Description

Recognizes a protein of 40kDa, identified as Thyroid transcription factor-1 (TTF-1) and Homeobox protein Nkx-2.1. NKX2.1 is a member of the NKx2 family of homeodomain transcription factors. It is expressed in epithelial cells of the thyroid gland and the lung. Nuclei from liver, stomach, pancreas, small intestine, colon, kidney, breast, skin, testes, pituitary, prostate, and adrenal glands are unreactive. Anti-NKX2.1 is useful in differentiating primary adenocarcinoma of the lung from metastatic carcinomas originating in the breast, mediastinal germ cell tumors, and malignant mesothelioma. It can also be used to differentiate small cell lung carcinoma from lymphoid infiltrates. Loss of NKX2.1 expression in non-small cell lung carcinoma has been associated with aggressive behavior of such neoplasms. NKX2.1 reactivity is also seen in thyroid malignancies.

Application Notes

Optimal dilution of the NKX2.1 antibody should be determined by the researcher.

Immunogen

A portion of amino acids 1-200 from human TTF-1 protein was used as the immunogen for the NKX2.1 antibody.

Storage

Aliquot the NKX2.1 antibody and store frozen at -20°C or colder. Avoid repeated freeze-thaw cycles.