


## NKX2.2 Antibody [clone RMNK2-2] (V3726)

Catalog No.	Formulation	Size
V3726-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V3726-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V3726SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug
V3726IHC-7ML	Prediluted in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide; *For IHC use only*	7 ml

**Bulk quote request**

<b>Species Reactivity</b>	Human
<b>Format</b>	Purified
<b>Clonality</b>	Monoclonal (mouse origin)
<b>Isotype</b>	Mouse IgG1, kappa
<b>Clone Name</b>	RMNK2-2
<b>Purity</b>	Protein G affinity chromatography
<b>UniProt</b>	O95096
<b>Gene ID</b>	4821
<b>Localization</b>	Nuclear
<b>Applications</b>	Immunohistochemistry (FFPE) : 1-2ug/ml for 30 min at RT Prediluted IHC Only Format : incubate for 30 min at RT (1)
<b>Limitations</b>	This NKX2.2 antibody is available for research use only.



IHC testing of FFPE Ewings sarcoma with NKX2.2 antibody (clone RMNK2-2). HIER: steam sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min.

## Description

Expression of NKX2.2 has been found in neuroendocrine tumors of the gut, making it a potential marker for the study of gastrointestinal neuroendocrine tumors. More recently, NKX2.2 protein was identified as a target of EWS-FLI-1, the fusion protein specific to Ewing sarcoma, and was shown to be differentially upregulated in Ewing sarcoma on the basis of array-based gene expression analysis. It acts as a valuable marker for Ewing sarcoma, with a sensitivity of 93% and a specificity of 89%, and aids in the differential diagnosis of small round cell tumors.

## Application Notes

Titration of the NKX2.2 antibody may be required for optimal performance.

1. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.

## Immunogen

A human partial recombinant protein corresponding to amino acids 1-119 was used as the immunogen for this NKX2.2 antibody.

## Storage

Store the NKX2.2 antibody at 2-8°C (with azide) or aliquot and store at -20°C or colder (without azide).