

NLRP3 Antibody / NALP3 (R30750)


Catalog No.	Formulation	Size
R30750	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug

Citations (1)

Flow Cytometry: 1-3ug/10^6 cells

This NLRP3 antibody is available for research use only.


	Oltations (1)	Bain quoto roquoe
Availability	1-3 business days	
Species Reactivity	Human, Mouse, Rat	
Format	Antigen affinity purified	
Clonality	Polyclonal (rabbit origin)	
Isotype	Rabbit Ig	
Purity	Antigen affinity	
Buffer	Lyophilized from 1X PBS with 2% Trehalose	
UniProt	Q96P20	
Localization	Nuclear, cytoplasmic, secreted	
Applications	Western Blot : 0.5-1ug/ml Immunohistochemistry (FFPE) : 2-5ug/ml	




Limitations

IHC staining of FFPE human squamous cell cervical carcinoma tissue with NLRP3 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.


Bulk quote request



IHC staining of FFPE human tonsil tissue with NLRP3 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.



Western blot testing of 1) human ThP-1, 2) rat PC-12, 3) mouse spleen and 4) mouse RAW264.7 cell lysate with NLRP3 antibody. Predicted molecular weight ~118 kDa.



Flow cytometry testing of fixed and permeabilized human ThP-1 cells with NLRP3 antibody at 1ug/million cells (blocked with goat sera); Red=cells alone, Green=isotype control, Blue= NLRP3 antibody.

Description

NOD-like receptor family pyrin domain containing 3, also known as CIAS1, CRYOPYRIN, NALP3 or PYPAF1, is a protein that in humans is encoded by the NLRP3 gene. The gene encodes a pyrin-like protein expressed predominantly in peripheral blood leukocytes. NLRP3 interacts with apoptosis-associated speck-like protein containing a CARD(ASC). The encoded protein may play a role in the regulation of inflammation and apoptosis. Mutation of the proteins nucleotide-binding domain reduced ATP binding, Caspase-1 activation, IL-1b production, cell death, macromolecular complex formation, self-association, and association with ASC. Consistent with an essential role for NLRP3 inflammasomes in antifungal immunity, Gross et al.showed that deficient mice are hypersusceptible to C. albicans infection. Activation of the NLRP3 inflammasome in response to virus or to RNA was dependent upon lysosomal maturation and reactive oxygen species production in human cells. The NLRP3 inflammasome senses obesity-associated danger signals and contributes to obesity-induced inflammation and insulin resistance.

Application Notes

The stated application concentrations are suggested starting points. Titration of the NLRP3 antibody may be required due to differences in protocols and secondary/substrate sensitivity.

Immunogen

An amino acid sequence from the N-terminus of human NLRP3 (RYLEDLEDVDLKKFKMHLED) was used as the immunogen for this NLRP3 antibody.

Storage

After reconstitution, the NLRP3 antibody can be stored for up to one month at 4oC. For long-term, aliquot and store at -20oC. Avoid repeated freezing and thawing.