


NMU Antibody / Neuromedin-U (RQ4097)


Catalog No.	Formulation	Size
RQ4097	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug

Bulk quote request


Availability	1-3 business days
Species Reactivity	Mouse, Rat
Format	Antigen affinity purified
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit IgG
Purity	Antigen affinity purified
Buffer	Lyophilized from 1X PBS with 2% Trehalose
UniProt	P48645
Applications	Western Blot : 0.5-1ug/ml Immunohistochemistry (FFPE) : 2-5ug/ml
Limitations	This NMU antibody is available for research use only.



IHC staining of FFPE mouse colon tissue with NMU antibody, HRP-secondary and DAB substrate. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.



IHC staining of FFPE rat colon tissue with NMU antibody, HRP-secondary and DAB substrate. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.



Western blot testing of 1) rat PC-12, 2) mouse kidney and 3) mouse Neuro-2a cell lysate with NMU antibody at 0.5ug/ml. Expected molecular weight: 20-25 kDa.

Description

NMU (Neuromedin-U) is a neuropeptide widely expressed in both the central nervous system and peripheral tissues. It is known for its conserved C-terminal pentapeptide motif, which is essential for receptor binding and biological activity. NMU exerts its physiological effects through interaction with two G protein-coupled receptors: NMUR1, primarily expressed in peripheral tissues, and NMUR2, which is predominantly found in the brain.

NMU plays a regulatory role in various biological functions, including energy balance, gastrointestinal activity, stress responses, and cardiovascular regulation. Its tissue-specific expression and receptor distribution make it a valuable target in studies of neuroendocrine signaling and metabolic control. Additionally, NMU expression is often evaluated in experimental models exploring neuropeptide function and receptor-ligand signaling pathways.

The **NMU antibody** is a reliable tool for detecting endogenous Neuromedin-U in applications such as western blot, immunohistochemistry, and immunofluorescence. Researchers use the NMU antibody from NSJ Bioreagents to monitor protein expression patterns, assess tissue localization, and study signaling dynamics in response to physiological or experimental stimuli. With high specificity and consistent performance, the NMU antibody supports detailed investigations into neuropeptide biology, receptor pathways, and regulatory peptide function.

Application Notes

Optimal dilution of the NMU antibody should be determined by the researcher.

Immunogen

Amino acids FRVDEEFQSPFASQSRGYFLFRPRN from the human protein were used as the immunogen for the NMU antibody.

Storage

After reconstitution, the NMU antibody can be stored for up to one month at 4°C. For long-term, aliquot and store at -20°C. Avoid repeated freezing and thawing.