

Nod1 Antibody (R31707)


Catalog No.	Formulation	Size
R31707	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug

Bulk quote request


Availability	1-3 business days
Species Reactivity	Human, Rat
Format	Antigen affinity purified
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit IgG
Purity	Antigen affinity
Buffer	Lyophilized from 1X PBS with 2.5% BSA, 0.025% sodium azide
Gene ID	10392
Applications	Western Blot : 0.5-1ug/ml IHC (FFPE) : 0.5-1ug/ml ICC (FFPE) : 0.5-1ug/ml
Limitations	This Nod1 antibody is available for research use only.




Western blot testing of Nod1 antibody and Lane 1: human A549; 2: rat heart lysate. Predicted/observed molecular weight: ~107 kDa.




IHC-P: Nod1 antibody testing of human placenta tissue. HIER: steam section in pH6 citrate buffer for 20 min and allow to cool prior to testing.



ICC testing of human SMMC-7721 cells with Nod1 antibody. HIER: steam section in pH6 citrate buffer for 20 min and allow to cool prior to testing.



ICC testing of human HeLa cells with Nod1 antibody. HIER: steam section in pH6 citrate buffer for 20 min and allow to cool prior to testing.



ICC testing of human A549 cells with Nod1 antibody. HIER: steam section in pH6 citrate buffer for 20 min and allow to cool prior to testing.

Description

Nucleotide-binding oligomerization domain-containing protein 1, also known as CARD4, is a protein receptor that in humans is encoded by the NOD1 gene. It is a member of NOD-like receptor protein family and is a close relative of Nod2. It recognizes bacterial molecules and stimulates an immune reaction. Nod1 contains a caspase recruitment domain (CARD). This gene is an intracellular pattern recognition receptor, which is similar in structure to resistant proteins of plants, and mediates innate and acquired immunity by recognizing bacterial molecules containing D-glutamyl-meso-diaminopimelic acid (iE-DAP) moiety. Additionally, it has been shown that Nod1 can sense cytosolic microbial products by monitoring the activation state of small Rho GTPases.

Application Notes

The stated application concentrations are suggested starting amounts. Titration of the Nod1 antibody may be required due to differences in protocols and secondary/substrate sensitivity.

Immunogen

Human partial recombinant protein (AA 1-160) was used as the immunogen for this Nod1 antibody.

Storage

After reconstitution, the Nod1 antibody can be stored for up to one month at 4oC. For long-term, aliquot and store at -20oC. Avoid repeated freezing and thawing.

