


## NONO Antibody (R34895)


Catalog No.	Formulation	Size
R34895-100UG	0.5 mg/ml in 1X TBS, pH7.3, with 0.5% BSA (US sourced) and 0.02% sodium azide	100 ug

[Bulk quote request](#)


<b>Availability</b>	1-3 business days
<b>Species Reactivity</b>	Human, Mouse
<b>Predicted Reactivity</b>	Dog, Rat
<b>Format</b>	Antigen affinity purified
<b>Host</b>	Goat
<b>Clonality</b>	Polyclonal (goat origin)
<b>Isotype</b>	Goat Ig
<b>Purity</b>	Antigen affinity
<b>UniProt</b>	Q15233
<b>Gene ID</b>	4841
<b>Localization</b>	Nuclear
<b>Applications</b>	Western Blot : 0.1-0.3ug/ml Immunohistochemistry (FFPE) : 3-6ug/ml ELISA (peptide) LOD : 1:16000
<b>Limitations</b>	This NONO antibody is available for research use only.




IHC staining of FFPE human spleen using NONO antibody at 3.8ug/ml. HIER: steamed with pH6 citrate buffer, AP-staining.



Western blot testing of mouse brain lysate with NONO antibody at 0.1ug/ml. Predicted molecular weight: ~54 kDa.



Western blot of HEK293 lysate overexpressing human NONO probed with NONO antibody (mock transfection in lane 1). Predicted molecular weight: ~54 kDa.



IHC staining of FFPE human testis using NONO antibody at 3.8ug/ml. HIER: steamed with pH6 citrate buffer, AP-staining.

## Description

Additional name(s) for this target protein: p54NRB, Non-POU domain containing octamer-binding

## Application Notes

Optimal dilution of the NONO antibody should be determined by the researcher.

## Immunogen

Amino acids NRAAPGAEFAPNK were used as the immunogen for this NONO antibody.

## Storage

Aliquot and store the NONO antibody at -20oC.