


NSE Antibody [clone NSEL1-1] (V7210)


Catalog No.	Formulation	Size
V7210-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V7210-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V7210SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

[Bulk quote request](#)

Species Reactivity	Human
Format	Purified
Host	Mouse
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG2b
Clone Name	NSEL1-1
Purity	Protein G affinity chromatography
Buffer	1X PBS, pH 7.4
UniProt	P09104
Gene ID	2026
Localization	Cytoplasmic
Applications	Western Blot : 1-2ug/ml Immunohistochemistry (FFPE) : 0.5-1ug/ml
Limitations	This NSE antibody is available for research use only.



IHC testing of FFPE human pancreas with NSE antibody (clone NSEL1-1). Staining of FFPE tissue requires boiling sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 min.



Western blot testing of human HeLa cell lysate with NSE antibody (clone NSEL1-1).
Predicted molecular weight ~47 kDa.

Description

NSE Antibody recognizes a protein of about 50kDa, which is identified as gamma-enolase/neuron specific enolase/enolase 2. Three isoenzymes of enolases are identified, alpha, beta and gamma. Alpha-isoform is expressed in most tissues, whereas beta-form is expressed predominantly in muscle tissue and gamma-enolase is found only in nervous tissue. These isoforms exist as both homodimers and heterodimers, and they play a role in converting phosphoglyceric acid to phosphoenolpyruvic acid in the glycolytic pathway. NSE is a useful marker to identify peripheral nerves and tumors of neuro-endocrine origins, such as pheochromocytomas. It is usually employed in combination with other markers such as Synaptophysin, Chromogranin A, and Neurofilament.

For detection of neuron-specific enolase (NSE), also known as gamma enolase, across tissue types, see our [NSE antibody](#).

Application Notes

The concentration stated for each application is a general starting point. Variations in protocols, secondaries and substrates may require the NSE antibody to be titrated up or down for optimal performance.

Immunogen

Amino acids 416-433 of human Neuron-specific Enolase were used as the immunogen for this NSE antibody.

Storage

Store the NSE antibody at 2-8°C (with azide) or aliquot and store at -20°C or colder (without azide).