


Nuclear Membrane Marker Antibody [clone NM97] (V2357)

Catalog No.	Formulation	Size
V2357-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V2357-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V2357SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

[Bulk quote request](#)

Species Reactivity	Human
Format	Purified
Host	Mouse
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG1, kappa
Clone Name	NM97
Purity	Protein G affinity chromatography
Buffer	1X PBS, pH 7.4
Gene ID	Unknown
Localization	Nuclear membrane
Applications	Flow Cytometry : 0.5-1ug/10 ⁶ cells Immunofluorescence : 0.5-1ug/ml
Limitations	This Nuclear membrane marker antibody is available for research use only.



SDS-PAGE Analysis of Purified, BSA-Free Nuclear Membrane Marker Antibody (clone NM97). Confirmation of Integrity and Purity of the Antibody.

Description

This monoclonal antibody is part of a panel of reagents which recognizes subcellular organelles or compartments of human cells. These markers may be useful in identification of these organelles in cells, tissues, and biochemical preparations. Clone NM97 antibody recognizes an antigenic marker associated with the nuclear membrane of human cells.

Application Notes

The concentration stated for each application is a general starting point. Variations in protocols, secondaries and substrates may require the Nuclear membrane marker antibody to be titered up or down for optimal performance.

Immunogen

Nuclei of myeloid leukemia biopsy cells were used as the immunogen for this Nuclear membrane marker antibody.

Storage

Store the Nuclear membrane marker antibody at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).