


Nucleoside diphosphate kinase B Antibody / NDKB / NME2 [clone NME2/6434] (V4518)


Catalog No.	Formulation	Size
V4518-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	100 ug
V4518-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	20 ug
V4518SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

Bulk quote request


Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG2, kappa
Clone Name	NME2/6434
Purity	Protein A/G affinity
UniProt	P22392
Localization	Cytoplasm, Nucleus
Applications	Western Blot : 1-2ug/ml Immunohistochemistry (FFPE) : 1-2ug/ml for 30 minutes at RT
Limitations	This Nucleoside diphosphate kinase B antibody is available for research use only.




IHC staining of FFPE human skin tissue with Nucleoside diphosphate kinase B antibody (clone NME2/6434). Inset: PBS used in place of primary Ab (secondary Ab negative control). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



SDS-PAGE analysis of purified, BSA-free Nucleoside diphosphate kinase B antibody (clone NME2/6434) as confirmation of integrity and purity.



Western blot testing of human 293T and MCF7 cell lysate with Nucleoside diphosphate kinase B antibody (clone NME2/6434). Predicted molecular weight ~17 kDa.



Analysis of a HuProt(TM) microarray containing more than 19,000 full-length human proteins using Nucleoside diphosphate kinase B antibody (clone NME2/6434). Z- and S-Score: The Z-score represents the strength of a signal that a monoclonal antibody (in combination with a fluorescently-tagged anti-IgG secondary antibody) produces when binding to a particular protein on the HuProt(TM) array. Z-scores are described in units of standard deviations (SDs) above the mean value of all signals generated on that array. If targets on HuProt(TM) are arranged in descending order of the Z-score, the S-score is the difference (also in units of SDs) between the Z-score. S-score therefore represents the relative target specificity of a mAb to its intended target. A mAb is considered to specific to its intended target, if the mAb has an S-score of at least 2.5. For example, if a mAb binds to protein X with a Z-score of 43 and to protein Y with a Z-score of 14, then the S-score for the binding of that mAb to protein X is equal to 29.

Description

The nm23 gene, a potential suppressor of metastasis, was originally identified by differential hybridization between two murine melanoma sub-lines, one with a high and the second with a low metastatic capacity. Highly metastatic sub-lines exhibit much lower levels of nm23 than less metastatic cells. Based on sequence analysis, nm23 appears highly related to nucleotide diphosphate kinases (NDP). In humans, NDP kinases A and B are identical to two isoforms of human nm23 homologs, namely nm23-H1 and H2, respectively. nm23-H2 is identical in sequence to PuF, a transcription factor that binds to nuclease hypersensitive elements at positions 142 to 115 of the human c-Myc promoter.

Application Notes

Optimal dilution of the Nucleoside diphosphate kinase B antibody should be determined by the researcher.

Immunogen

Recombinant full-length human protein was used as the immunogen for the Nucleoside diphosphate kinase B antibody.

Storage

Aliquot the Nucleoside diphosphate kinase B antibody and store frozen at -20°C or colder. Avoid repeated freeze-thaw cycles.

