


# NUR77 Antibody (R30761)

| Catalog No. | Formulation   | Size   |
|-------------|---|--------|
| R30761      | 0.5mg/ml if reconstituted with 0.2ml sterile DI water | 100 ug |

## **Bulk quote request**

| Availability       | 1-3 business days  |
|--------------------|--|
| Species Reactivity | Human, Mouse, Rat  |
| Format             | Antigen affinity purified  |
| Clonality          | Polyclonal (rabbit origin)   |
| Isotype            | Rabbit Ig  |
| Purity             | Antigen affinity   |
| Buffer             | Lyophilized from 1X PBS with 2.5% BSA and 0.025% sodium azide/thimerosal |
| UniProt            | P22736   |
| Applications       | Western Blot : 0.5-1ug/ml  |
| Limitations        | This NUR77 antibody is available for research use only.                  |



## **Description**

Nuclear receptor subfamily 4 group A member 1, also called NAK1, GFRP1, TR3, NUR77 or NGFIB, is a protein that in humans is encoded by the NR4A1 gene, and is a member of the Nur nuclear receptor family of intracellular transcription factors. NR4A1 is involved in cell cycle mediation, inflammation and apoptosis. It plays a key role in mediating inflammatory responses in macrophages. In addition, subcellular localization of the protein appears to play a key role in the survival and death of cells. Nr4a1 was overexpressed in Wnt1 -transformed mouse mammary cells. It was also induced by lithium, a Wnt1 mimic, and the promoter was activated by lithium and beta-catenin, a Wnt1 downstream effector. In contrast, human NR4A1 was not upregulated by beta-catenin, indicating that this gene is regulated differently in human and mouse cells. Adenoviral expression of Nr4a1 induced genes involved in gluconeogenesis, stimulated

glucose production both in vitro and in vivo, and raised blood glucose levels.

#### **Application Notes**

The stated application concentrations are suggested starting points. Titration of the NUR77 antibody may be required due to differences in protocols and secondary/substrate sensitivity.

## **Immunogen**

An amino acid sequence from the N-terminus of NUR77 alpha/NR4A1 (ASFKFEDFQVYGCY) was used as the immunogen for this NUR77 antibody (100% homologous in human, mouse and rat).

#### **Storage**

After reconstitution, the NUR77 antibody can be stored for up to one month at 4oC. For long-term, aliquot and store at -20oC. Avoid repeated freezing and thawing.

#### **Alternate Names**

NAK1, GFRP1, TR3, NGFIB