


Olr1 Antibody / Lox-1 (RQ6063)

Catalog No.	Formulation	Size
RQ6063	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug

[Bulk quote request](#)

Availability	1-3 business days
Species Reactivity	Mouse
Format	Antigen affinity purified
Host	Rabbit
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit IgG
Purity	Affinity purified
Buffer	Lyophilized from 1X PBS with 2% Trehalose and 0.025% sodium azide
UniProt	Q9EQ09
Applications	Western Blot : 0.5-1ug/ml Direct ELISA : 0.1-0.5ug/ml
Limitations	This Olr1 antibody is available for research use only.



Western blot testing of mouse 1) brain, 2) lung, 3) liver, 4) kidney, 5) testis, 6) smooth muscle, 7) SP2/0 and 8) Neuro-2a cell lysate with Olr1 antibody. Predicted molecular weight: pro-form 35-50 kDa, mature form ~31 kDa.

Description

OLR1 (oxidized low density lipoprotein (lectin-like) receptor 1) also called CLEC8A, LOX-1, SCARE1, is a receptor protein which belongs to the C-type lectin superfamily. The OLR1 gene encodes a cell-surface endocytosis receptor for oxidized low density lipoprotein (OxLDL). This gene is mapped on 12p13.2. Incubation of the cells with LDL had no effect on LOX1 expression, but incubation with OxLDL resulted in a dose-dependent increase in LOX1 mRNA and protein expression; however, very high concentrations of OxLDL caused a decrease in OxLDL expression, perhaps indicating toxic effects on endothelial cells. LOX1 was also expressed in macrophages, but not in vascular smooth muscle cells. The findings

suggested a role for LOX1 in the pathophysiology of atherosclerotic cardiovascular disease. LOX1 expression was detected in all choroidal neovascular membranes, regardless of structure, whereas there was no evidence of LOX1 within the posterior segments of normal eyes. LOX1 plays an active role in the pathogenesis of choroidal neovascularization, especially in ARMD.

Application Notes

Optimal dilution of the Olr1 antibody should be determined by the researcher.

Immunogen

Recombinant mouse protein (amino acids Q55-I363) was used as the immunogen for the Olr1 antibody.

Storage

After reconstitution, the Olr1 antibody can be stored for up to one month at 4°C. For long-term, aliquot and store at -20°C. Avoid repeated freezing and thawing.