


Opa-interacting protein 4 Antibody / OIP-4 / PRAME [clone PRAME/9189] (V5572)


Catalog No.	Formulation	Size
V5572-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	100 ug
V5572-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	20 ug
V5572SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

Bulk quote request

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Host	Mouse
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG2b, kappa
Clone Name	PRAME/9189
Purity	Protein A/G affinity
UniProt	P78395
Localization	Nucleus, Cytoplasm, Cell membrane
Applications	Immunohistochemistry (FFPE) : 1-2ug/ml
Limitations	This Opa-interacting protein 4 antibody is available for research use only.



IHC staining of FFPE human melanoma tissue with Opa-interacting protein 4 antibody (clone PRAME/9189). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



Description

Functions as a transcriptional repressor, inhibiting the signaling of retinoic acid through the retinoic acid receptors RARA, RARB and RARG. Prevents retinoic acid-induced cell proliferation arrest, differentiation and apoptosis. This gene encodes an antigen that is preferentially expressed in human melanomas and that is recognized by cytolytic T lymphocytes. It is not expressed in normal tissues, except testis. The encoded protein acts as a repressor of retinoic acid receptor, and likely confers a growth advantage to cancer cells via this function. Alternative splicing results in multiple transcript variants.

Application Notes

Optimal dilution of the PRAME antibody should be determined by the researcher.

Immunogen

A recombinant fragment (within amino acids 100-250) of human PRAME protein was used as the immunogen for the Opa-interacting protein 4 antibody.

Storage

Aliquot the PRAME antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.