

P-Cadherin Antibody / CDH3 [clone 6A9] (V3397)


Catalog No.	Formulation	Size
V3397-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V3397-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V3397SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

 [Citations \(21\)](#)
[Bulk quote request](#)

Species Reactivity	Human
Format	Purified
Host	Mouse
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG1, kappa
Clone Name	6A9
Purity	Protein G affinity chromatography
Buffer	1X PBS, pH 7.4
UniProt	P22223
Localization	Cell surface, cytoplasmic
Applications	Flow Cytometry : 0.5-1ug/10 ⁶ cells Immunofluorescence : 1-2ug/ml Western Blot : 0.5-1ug/ml
Limitations	This P-Cadherin antibody is available for research use only.



Western blot testing of human A431 cell lysate with P-Cadherin antibody (clone 6A9).
Expected molecular weight: ~91 kDa (unmodified), 100~130 kDa (glycosylated).



SDS-PAGE Analysis of Purified, BSA-Free P-Cadherin Antibody (clone 6A9).
Confirmation of Integrity and Purity of the Antibody.

Description

Recognizes a protein of 116kDa, identified as P-Cadherin, or Cadherin 3 or CDH3. It is a calcium-dependent cell-cell adhesion glycoprotein comprised of five extracellular cadherin repeats, a transmembrane region and a highly conserved cytoplasmic tail. This gene is located in a six-cadherin cluster in a region on the long arm of chromosome 16 that is involved in loss of heterozygosity events in breast and prostate cancer. In addition, aberrant expression of this protein is observed in cervical adenocarcinomas. Mutations in this gene have been associated with congenital hypotrichosis with juvenile macular dystrophy.

Application Notes

The concentration stated for each application is a general starting point. Variations in protocols, secondaries and substrates may require the P-Cadherin antibody to be titrated up or down for optimal performance.

Immunogen

Membrane from trypsinized A431 cells was used as the immunogen for this P-Cadherin antibody.

Storage

Store the P-Cadherin antibody at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).