


p-ERBB2 Antibody (pY1196) (F48699)


Catalog No.	Formulation	Size
F48699-0.4ML	In 1X PBS, pH 7.4, with 0.09% sodium azide	0.4 ml
F48699-0.08ML	In 1X PBS, pH 7.4, with 0.09% sodium azide	0.08 ml

[Bulk quote request](#)

Availability	1-3 business days
Species Reactivity	Human
Format	Antigen affinity purified
Host	Rabbit
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit Ig
Purity	Antigen affinity
UniProt	P04626
Applications	Western Blot : 1:4000 Immunofluorescence : 1:100 Dot Blot : 1:500
Limitations	This p-ERBB2 antibody is available for research use only.




Western blot analysis of extracts from A431 cells, untreated or treated with EGF (100ng/ml) using p-ERBB2 antibody (left) or nonphos Ab (right)



Fluorescent confocal image of MCF7 cells stained with p-ERBB2 antibody. MCF7 cells were fixed with 4% PFA (20 min), permeabilized with Triton X-100 (0.2%, 30 min). Cells were then incubated with primary Ab (1:100, 2 h at room temperature). For secondary Ab, Alexa Fluor 488 conjugated donkey anti-rabbit Ab (green) was used (1:1000, 1h). Nuclei were counterstained with Hoechst 33342 (blue) (10 ug/ml, 5 min). Note the highly specific localization to the plasma membrane and cytoplasm.

Dot blot analysis of p-ERBB2 antibody. 50ng of phos-peptide or nonphos-peptide per dot were spotted.



Description

This gene encodes a member of the epidermal growth factor (EGF) receptor family of receptor tyrosine kinases. This protein has no ligand binding domain of its own and therefore cannot bind growth factors. However, it does bind tightly to other ligand-bound EGF receptor family members to form a heterodimer, stabilizing ligand binding and enhancing kinase-mediated activation of downstream signalling pathways, such as those involving mitogen-activated protein kinase and phosphatidylinositol-3 kinase. Allelic variations at amino acid positions 654 and 655 of isoform a (positions 624 and 625 of isoform b) have been reported, with the most common allele, Ile654/Ile655, shown here. Amplification and/or overexpression of this gene has been reported in numerous cancers, including breast and ovarian tumors. Alternative splicing results in several additional transcript variants, some encoding different isoforms and others that have not been fully characterized.

Application Notes

Titration of the p-ERBB2 antibody may be required due to differences in protocols and secondary/substrate sensitivity.

Immunogen

This p-ERBB2 antibody was produced from rabbits immunized with a KLH conjugated synthetic phosphopeptide corresponding to amino acid residues surrounding pY1196 of human ERBB2.

Storage

Aliquot the p-ERBB2 antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.