


# P-glycoprotein Antibody / MDR1 / ABCB1 [clone MDR1/8962R] (V5207)

Catalog No.	Formulation	Size
V5207-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	100 ug
V5207-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	20 ug
V5207SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

### Recombinant RABBIT MONOCLONAL

### **Bulk quote request**

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Clonality	Recombinant Rabbit Monoclonal
Isotype	Rabbit IgG, kappa
Clone Name	MDR1/8962R
Purity	Protein A/G affinity
UniProt	P08183
Localization	Membrane
Applications	Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT
Limitations	This P-glycoprotein antibody is available for research use only.



IHC staining of FFPE human adrenal gland tissue with MDR1 antibody (clone MDR1/8962R). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.

## **Description**

Cells selected for resistance to a single cytotoxic drug may become crossresistant to a broad range of drugs with different structures and cellular targets. This phenomenon is called multiple drug resistance (MDR). MDR proteins (Mdrs) are members of a highly conserved superfamily of ATP-binding cassette transport proteins. Mdr-1 is an apical transmembrane protein that is an integral part of the blood-brain barrier and functions as a drug-transport pump

transporting a variety of drugs from the brain back into the blood. The Mdr-1 gene is known as ABCB1 and is located on human chromosome 7q21.12. The mouse homolog of Mdr-1 is known as Mdr-3. Interestingly, a murine protein by the name of Mdr-1 exists and is encoded by the murine Abcb1b gene, but it is not homologous with human Mdr-1.

### **Application Notes**

Optimal dilution of the P-glycoprotein antibody should be determined by the researcher.

### **Immunogen**

A recombinant partial protein sequence (within amino acids 500-700) from the human protein was used as the immunogen for the P-glycoprotein antibody.

### **Storage**

Aliquot the P-glycoprotein antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.