
P Glycoprotein Antibody (RQ4527)


Catalog No.	Formulation	Size
RQ4527	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug

Bulk quote request


Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Host	Rabbit
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit IgG
Purity	Protein A affinity
Buffer	Lyophilized from 1X PBS with 2% Trehalose and 0.025% sodium azide
UniProt	P08183
Localization	Cell membrane
Applications	Western blot: 0.5-1ug/ml Immunofluorescence (FFPE): 2-4ug/ml Flow cytometry: 1-3ug/million cells
Limitations	This P Glycoprotein antibody is available for research use only.



Immunofluorescent staining of FFPE human U-2 OS cells with P Glycoprotein antibody (green) and DAPI nuclear stain (blue). HIER: steam section in pH6 citrate buffer for 20 min.



Flow cytometry testing of human U-2 OS cells with P Glycoprotein antibody at 1ug/million cells (blocked with goat sera); Red=cells alone, Green=isotype control, Blue= P Glycoprotein antibody.



Western blot testing of human 1) THP-1 and 2) A375 cell lysate with P Glycoprotein antibody at 0.5ug/ml. Expected molecular weight: 141-180 kDa depending on glycosylation level.

Description

P-GP, also called ABCB1 or PGY1, is a glycoprotein that in humans is encoded by the ABCB1 gene. It is mapped to 7q21.12. P-GP is a well-characterized ABC-transporter (which transports a wide variety of substrates across extra- and intracellular membranes) of the MDR/TAP subfamily. It is an important protein of the cell membrane that pumps many foreign substances out of cells. More formally, it is an ATP-dependent drug efflux pump with broad substrate specificity. P-GP is an ATP-dependent drug efflux pump forxenobiotic compounds with broad substrate specificity. It is responsible for decreased drug accumulation in multidrug-resistant cells and often mediates the development of resistance to anticancer drugs. This protein also functions as a transporter in the blood-brain barrier.

Application Notes

Optimal dilution of the P Glycoprotein antibody should be determined by the researcher.

Immunogen

Amino acids QAQDRKLSTKEALDESIPPVSFWRIMKLNLTEWPY were used as the immunogen for the P Glycoprotein antibody antibody.

Storage

After reconstitution, the P Glycoprotein antibody can be stored for up to one month at 4oC. For long-term, aliquot and store at -20oC. Avoid repeated freezing and thawing.

Ordering: Phone: 858.663.9055 | Fax: 1.267.821.0800 | Email: info@nsjbio.com
Copyright © NSJ Bioreagents. All rights reserved.