


p-NANOG Antibody (pS71) (F48601)


Catalog No.	Formulation	Size
F48601-0.4ML	In 1X PBS, pH 7.4, with 0.09% sodium azide	0.4 ml
F48601-0.08ML	In 1X PBS, pH 7.4, with 0.09% sodium azide	0.08 ml

[Bulk quote request](#)

Availability	1-3 business days
Species Reactivity	Human
Predicted Reactivity	Primate
Format	Antigen affinity purified
Host	Rabbit
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit Ig
Purity	Antigen affinity
UniProt	Q9H9S0
Applications	Immunofluorescence : 1:10-1:50 Dot Blot : 1:500
Limitations	This p-NANOG antibody is available for research use only.



Fluorescent confocal image of SY5Y cells stained with p-NANOG antibody at 1:50. Immunoreactivity is localized very specifically to the nuclei.



Dot blot analysis of p-NANOG antibody. 50ng of phos-peptide or nonphos-peptide per dot were spotted.

Description

NANOG is a transcription regulator involved in inner cell mass and embryonic stem (ES) cells proliferation and self-renewal. It imposes pluripotency on ES cells and prevents their differentiation towards extraembryonic endoderm and trophoderm lineages. This protein blocks bone morphogenetic protein-induced mesoderm differentiation of ES cells by physically interacting with SMAD1 and interfering with the recruitment of coactivators to the active SMAD transcriptional complexes. NANOG acts as a transcriptional activator or repressor. It binds optimally to the DNA consensus sequence 5'-[CG][GA][CG]C[GC]ATTAN[GC]-3'. When overexpressed, this protein promotes cells to enter into S phase and proliferation.

Application Notes

Titration of the p-NANOG antibody may be required due to differences in protocols and secondary/substrate sensitivity.

Immunogen

This p-NANOG antibody was produced from rabbits immunized with a KLH conjugated synthetic phosphopeptide corresponding to amino acid residues surrounding pS71 of human Nanog.

Storage

Aliquot the p-NANOG antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.