

## p120 Catenin Antibody (pTyr96) / CTNND1 / Catenin delta 1 [clone 25a] (V7701)

Catalog No.	Formulation	Size
V7701-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V7701-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V7701SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

### Bulk quote request

Availability	1-3 business days
Species Reactivity	Human, Mouse, Rat
Format	Purified
Host	Mouse
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG1, kappa
Clone Name	25a
Purity	Protein G affinity chromatography
UniProt	O60716
Applications	Flow Cytometry : 1-2ug/10 <sup>6</sup> cells in 0.1ml Immunofluorescence : 1-2ug/ml
Limitations	This p120 Catenin antibody is available for research use only.



### Description

The membrane associated protein pp120 Src substrate (p120 Catenin, p120cas) was identified as a tyrosine kinase substrate that is phosphorylated in Src transformed cells or in response to growth factor stimulation. It shares structural similarity with the *Drosophila* Armadillo protein and the vertebrate beta-catenin and gamma-catenin proteins. Its characteristic Arm domain that is composed of 42-amino acid motif repeats evidences this similarity. In the cell, p120 Catenin is localized to the E-Cadherin/catenins cell adhesion complex. Like beta- and gamma-catenin, p120 Catenin directly associates with the cytoplasmic C-terminus of E-Cadherin via its Arm domain and may similarly interact with other Cadherins. It exists as four isoforms that range in size from 90-115kDa. Expression of these isoforms is heterogeneous in human carcinomas, suggesting that altered pp120 expression contributes to malignancy due to loss of functional cell adhesions. Multiple tyrosine residues (Y96, Y112, Y228, Y280, Y257, Y291, Y296, and Y302) in p120 Catenin are phosphorylated by Src and these phosphorylations may facilitate interaction with PTP1C/SHP-1 in response to EGF stimulation. Thus, p120 Catenin is an Arm domain protein that interacts with both cell adhesion molecules, such as cadherins and cell signaling molecules, such as PTP1C.

## Application Notes

Optimal dilution of the p120 Catenin antibody should be determined by the researcher.

## Immunogen

The amino acid sequence surrounding phosphorylated Tyr96 was used as the immunogen for the p120 Catenin antibody.

## Storage

Store the p120 Catenin antibody at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).