


p14ARF Antibody / CDKN2A [clone 4C6/4] (V3807)


Catalog No.	Formulation	Size
V3807-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V3807-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V3807SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

Bulk quote request


Availability	1-3 business days
Species Reactivity	Human, Rat
Format	Purified
Host	Mouse
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG2a, kappa
Clone Name	4C6/4
Purity	Protein G affinity chromatography
UniProt	P42771
Localization	Nuclear, cytoplasmic
Applications	Flow Cytometry : 1-2ug/10 ⁶ cells Immunofluorescence : 1-2ug/ml Western Blot : 1-2ug/ml Immunohistochemistry (FFPE) : 1-2ug/ml for 30 min at RT
Limitations	This p14ARF antibody is available for research use only.



IHC testing of FFPE human brain with p14ARF antibody (clone 4C6/4). HIER: boil tissue sections in pH6, 10mM citrate buffer, for 10-20 min followed by cooling at RT for 20 min.



IHC testing of FFPE rat brain with p14ARF antibody (clone 4C6/4). HIER: boil tissue sections in pH6, 10mM citrate buffer, for 10-20 min followed by cooling at RT for 20 min.



SDS-PAGE analysis of purified, BSA-free p14ARF antibody (clone 4C6/4) as confirmation of integrity and purity.

Description

The progression of cells through the cell cycle is regulated by a family of proteins designated cyclin-dependent kinases (Cdks). Sequential activation of individual members of this family and their consequent phosphorylation of critical substrates promotes orderly progression through the cell cycle. Multiple proteins are encoded by the tumor suppressor gene CDKN2A (MTS1/ p16INK4a) via translation of alternate reading frames, resulting in the production of the p19ARF protein in mice and the p14ARF protein in humans. p14ARF induces an increase in MDM2 and p21 levels and leads to cell cycle arrest in both G1 and G2/M. p14ARF is negatively regulated by p53 and is known to bind directly to MDM2. CDKN2A also encodes the mitotic protein p16, which binds to and inhibits the Cdk4/cyclin D complex.

Application Notes

Optimal dilution of the p14ARF antibody should be determined by the researcher.

Immunogen

Full-length recombinant human protein used as the immunogen for the p14ARF antibody.

Storage

Store the p14ARF antibody at 2-8°C (with azide) or aliquot and store at -20°C or colder (without azide).