


# p16INK4a Antibody / CDKN2A [clone CDKN2A/4499] (V8604)


| Catalog No.    | Formulation  | Size   |
|----------------|--|--------|
| V8604-100UG    | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide | 100 ug |
| V8604-20UG     | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide | 20 ug  |
| V8604SAF-100UG | 1 mg/ml in 1X PBS; BSA free, sodium azide free                             | 100 ug |

## **Bulk quote request**


| Availability       | 1-3 business days  |
|--------------------|--|
| Species Reactivity | Human  |
| Format             | Purified   |
| Clonality          | Monoclonal (mouse origin)                                  |
| Isotype            | Mouse IgG2a, kappa   |
| Clone Name         | CDKN2A/4499  |
| Purity             | Protein G affinity chromatography                          |
| UniProt            | P42771   |
| Localization       | Nuclear and cytoplasmic                                    |
| Applications       | Immunohistochemistry (FFPE): 1-2ug/ml for 30 minutes at RT |
| Limitations        | This p16INK4a antibody is available for research use only. |



IHC staining of FFPE human cervix with p16INK4a antibody (clone CDKN2A/4499). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



IHC staining of FFPE human cervix with p16INK4a antibody (clone CDKN2A/4499). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



SDS-PAGE analysis of purified, BSA-free p16INK4a antibody (clone CDKN2A/4499) as confirmation of integrity and purity.

## **Description**

p16INK4a is a tumor suppressor protein. It is a specificinhibitor of cdk4/cdk6, and a tumorsuppressor involved inthe pathogenesis of a variety of malignancies. Recentanalyses of the p16INK4a gene revealed homozygousdeletions, nonsense, missense, or frameshift mutations inseveral human cancers. Although the frequency of p16INK4aabnormalities is higher in tumor derived cell lines than inunselected primary tumors, significant subsets of clinicalcases with aberrant p16INK4a gene have been reportedamong melanomas, gliomas, esophageal, pancreatic, lung,and urinary bladdercarcinomas, and some types of leukemia. Expression of p16INK4a (p16 positive) is highly correlated with human papilloma virus (HPV) infection in head and necksquamous cell carcinomas (HNSCC). p16 status is animportant prognostic indicator in HNSCC and the p16positive/HPV16 negative group is likely a distinct subgrouplacking any HPV genotype.

### **Application Notes**

Optimal dilution of the p16INK4a antibody should be determined by the researcher.

#### **Immunogen**

Recombinant full-length human protein was used as the immunogen for the p16INK4a antibody.

#### **Storage**

Store the p16INK4a antibody at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).