


p27Kip1 Antibody [clone KIP27-1] (V7048)


Catalog No.	Formulation	Size
V7048-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V7048-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V7048SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug
V7048IHC-7ML	Prediluted in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide; *For IHC use only*	7 ml

[Bulk quote request](#)


Availability	1-3 business days
Species Reactivity	Human, Mouse and Rat
Format	Purified
Host	Mouse
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG1, kappa
Clone Name	KIP27-1
Purity	Protein G affinity chromatography
UniProt	P46527
Localization	Nuclear
Applications	Flow Cytometry : 1-2ug/million cells in 0.1ml Immunofluorescence : 1-2ug/ml Immunohistochemistry (FFPE) : 0.25-0.5ug/ml for 30 min at RT Prediluted IHC Only Format : incubate for 30 min at RT (1)
Limitations	This p27Kip1 antibody is available for research use only.




IHC testing of FFPE human prostate cancer and p27Kip1 antibody (clone KIP27-1). Staining of formalin-fixed tissues requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 min.




IHC testing of FFPE human colon carcinoma and p27Kip1 antibody (clone KIP27-1). Staining of formalin-fixed tissues requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 min.



IHC testing of FFPE rat colon tissue and p27Kip1 antibody (clone KIP27-1). Staining of formalin-fixed tissues requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 min.



FACS testing of Jurkat cells and Alexa Fluor 488-labeled p27Kip1 antibody.



Immunofluorescent staining of PFA-fixed human MCF7 cells with p27Kip1 antibody (clone KIP27-1, green) and Phalloidin (red).

Description

This mAb recognizes a 27kDa protein, identified as the p27Kip1, a cell cycle regulatory mitotic inhibitor. It is highly specific and shows no cross-reaction with other related mitotic inhibitors. p27Kip1 functions as a negative regulator of G1 progression and has been proposed to function as a possible mediator of TGF- induced G1 arrest. p27Kip1 is a candidate tumor suppressor gene.

Application Notes

Optimal dilution of the p27Kip1 antibody to be determined by the researcher.

1. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.

Immunogen

Recombinant human protein was used as the immunogen for the p27Kip1 antibody.

Storage

Store the p27Kip1 antibody at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).