
p53 Antibody (V3343)

Catalog No.	Formulation	Size
V3343-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V3343-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V3343SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

Bulk quote request

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Host	Rabbit
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit IgG
Purity	Protein A affinity chromatography
UniProt	P04637
Localization	Predominantly nuclear
Applications	Flow cytometry: 0.5-1ug/10^6 cells Immunofluorescence: 0.5-1ug/ml Western blot: 0.5-1ug/ml Immunohistochemistry (FFPE): 0.25-0.5ug/ml for 30 min at RT
Limitations	This p53 antibody is available for research use only.



IHC testing of FFPE human colon carcinoma with p53 antibody. Required HIER: boil tissue sections in 10mM citrate buffer, pH 6, for 10-20 min.

Description

Recognizes a 53kDa protein, which is identified as p53 suppressor gene product. It reacts with the mutant as well as the wild form of p53. p53 is a tumor suppressor gene expressed in a wide variety of tissue types and is involved in regulating cell growth, replication, and apoptosis. It binds to MDM2, SV40 T antigen and human papilloma virus E6 protein. Positive nuclear staining with p53 antibody has been reported to be a negative prognostic factor in breast carcinoma, lung carcinoma, colorectal, and urothelial carcinoma. Anti-p53 positivity has also been used to differentiate uterine serous carcinoma from endometrioid carcinoma as well as to detect intratubular germ cell neoplasia. Mutations involving p53 are found in a wide variety of malignant tumors, including breast, ovarian, bladder, colon, lung, and melanoma.

Researchers studying p53 expression and TP53 signaling can explore our Protein Microarray Validated p53 Antibody page for additional reagents targeting this key tumor suppressor in cancer, DNA damage response and cell cycle research.

Application Notes

The optimal dilution of the p53 antibody for each application should be determined by the researcher.

Immunogen

Recombinant human p53 protein was used as the immunogen for this p53 antibody.

Storage

Store the p53 antibody at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).

Copyright © NSJ Bioreagents. All rights reserved.