


p73 Antibody [clone P73/2531] (V8089)


Catalog No.	Formulation	Size
V8089-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V8089-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V8089SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

Bulk quote request

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG2b, kappa
Clone Name	P73/2531
Purity	Protein G affinity chromatography
UniProt	O15350
Localization	Nuclear
Applications	Immunohistochemistry (FFPE) : 1-2ug/ml
Limitations	This p73 antibody is available for research use only.



IHC staining of FFPE human skin with p73 antibody. HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



SDS-PAGE analysis of purified, BSA-free p73 antibody as confirmation of integrity and purity.

Description

The p53 gene is a widely studied anti-oncogene, or tumor suppressor gene. The p53 gene product can act as a negative regulator of cell growth in response to DNA damage. Mutations and allelic loss of the p53 gene have been associated with malignant transformation in a wide variety of human tumors. p53 shares considerable sequence similarity with p73, a gene that maps to a region in chromosome 1 that is frequently deleted in neuroblastomas. However, p73 does not appear to be activated by DNA damaging agents.

Application Notes

Optimal dilution of the p73 antibody should be determined by the researcher.

Immunogen

A recombinant full-length human protein was used as the immunogen for this p73 antibody.

Storage

Store the p73 antibody at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).